

Tuesday, February 2, 2021

AM Session

Speaker: Richard Rimanyi, University of North Carolina at
Chapel Hill

Teaching Assistants: David Oetjen, Yiyan Shou, Xiping Zhang,
Rahul Singh, Changjian Su

**BASIC NOTIONS IN COTANGENT SCHUBERT CALCULUS
INTRO WORKSHOP ON COMBINATORIAL ALGEBARIC GEOMETRY
ICERM 2021
PROBLEMS**

R. RIMÁNYI

These problems use the notations, and refer to notions from my lecture “*Basic notions in cotangent Schubert Calculus*” at CIRM 2021. Lecture notes are available upon request. The meaning and relevance of the statements made in these problems are also explained in that lecture.

Problem 1 (on $H_T^*(\mathbb{P}^1)$)

Consider the ring homomorphism

$$\begin{aligned} \text{Loc} : \mathbb{Z}[t, z_1, z_2] &\rightarrow \mathbb{Z}[z_1, z_2] \oplus \mathbb{Z}[z_1, z_2] \\ f(t, z_1, z_2) &\mapsto (f(z_1, z_1, z_2), f(z_2, z_1, z_2)). \end{aligned}$$

Prove the following two characterizations of the image (range) of Loc:

$$\begin{aligned} \text{Im}(\text{Loc}) &= \{(f_1(z_1, z_2), f_2(z_1, z_2)) \in \mathbb{Z}[z_1, z_2] \oplus \mathbb{Z}[z_1, z_2] : f_1(u, u) = f_2(u, u)\}, \\ \text{Im}(\text{Loc}) &= \{(f_1(z_1, z_2), f_2(z_1, z_2)) \in \mathbb{Z}[z_1, z_2] \oplus \mathbb{Z}[z_1, z_2] : (z_1 - z_2) \mid (f_1 - f_2)\}. \end{aligned}$$

Problem 2 (on equivariant Schubert classes in $H_T^*(\mathbb{P}^{n-1})$)

Let $j \leq n$ be non-negative integers. Invent (that is, give a formula for) a polynomial $f(t, z_1, z_2, \dots, z_n)$ such that

- f is of homogeneous degree $n - j$ (where $\deg t = \deg z_i = 1$);
- $f(z_j, z_1, z_2, \dots, z_n) = \prod_{i=j+1}^n (z_i - z_j)$;
- $f(z_i, z_1, z_2, \dots, z_n) = 0$ if $j < i \leq n$.

Problem 3 (on equivariant CSM classes in $H_T^*(\mathbb{P}^{n-1})$)

Let $j \leq n$ be non-negative integers. Invent a polynomial $f(t, z_1, z_2, \dots, z_n, \hbar)$ such that

- f is of homogeneous degree $n - 1$ (where $\deg t = \deg z_i = \deg \hbar = 1$);
- $f(z_j, z_1, z_2, \dots, z_n) = \prod_{i=1}^{j-1} (z_i - z_j + \hbar) \prod_{i=j+1}^n (z_i - z_j)$;
- $f(z_i, z_1, z_2, \dots, z_n) = 0$ if $j < i \leq n$;
- $f(z_i, z_1, z_2, \dots, z_n)$ is divisible by \hbar for $i < j$;
- $f(z_i, z_1, z_2, \dots, z_n)$ is divisible by $\prod_{s=1}^{i-1} (z_s - z_i + \hbar)$.

Problem 4 (on the MacPherson property of CSM classes)

Consider the polynomial f you defined in Problem 3, and let us call it $f_{j,n}$. Define $F_n = \sum_{j=1}^n f_{j,n}$. Show that $F_n|_{t=z_i}$ is a product of linear factors, for all n and i .

Problem 5 (on equivariant Littlewood-Richardson coefficients on \mathbb{P}^1)

In the lecture we saw that in $H_T^*(\mathbb{P}^1)$ we have

$$\begin{aligned} [\overline{\Omega}_1] &= (z_2 - z_1, 0), \\ [\overline{\Omega}_2] &= (1, 1). \end{aligned}$$

Calculate the products $[\overline{\Omega}_i] \cdot [\overline{\Omega}_j]$ as $\mathbb{Z}[z_1, z_2]$ -linear combinations of $[\overline{\Omega}_1]$ and $[\overline{\Omega}_2]$.

Problem 6 (on CSM versions of equivariant Littlewood-Richardson coefficients on \mathbb{P}^1)

In the lecture we saw that in $H_T^*(\mathbb{P}^1)$ we have

$$\begin{aligned} c^{\text{sm}}(\Omega_1) &= (z_2 - z_1, 0), \\ c^{\text{sm}}(\Omega_2) &= (\hbar, z_1 - z_2 + \hbar). \end{aligned}$$

Calculate the products $c^{\text{sm}}(\Omega_i) \cdot c^{\text{sm}}(\Omega_j)$ as $\mathbb{Z}[z_1, z_2, \hbar]$ -linear combinations of $c^{\text{sm}}(\Omega_1)$ and $c^{\text{sm}}(\Omega_2)$.

Problem 7 (on the R -matrix property on CSM classes)

In the lecture we claimed that

$$(1) \quad \begin{pmatrix} c^{\text{sm}}(\Omega_{\emptyset}^{\text{opposite}}) \\ c^{\text{sm}}(\Omega_1^{\text{opposite}}) \\ c^{\text{sm}}(\Omega_2^{\text{opposite}}) \\ c^{\text{sm}}(\Omega_{12}^{\text{opposite}}) \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \frac{z_1 - z_2}{z_1 - z_2 + \hbar} & \frac{\hbar}{z_1 - z_2 + \hbar} & 0 \\ 0 & \frac{\hbar}{z_1 - z_2 + \hbar} & \frac{z_1 - z_2}{z_1 - z_2 + \hbar} & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} c^{\text{sm}}(\Omega_{\emptyset}) \\ c^{\text{sm}}(\Omega_1) \\ c^{\text{sm}}(\Omega_2) \\ c^{\text{sm}}(\Omega_{12}) \end{pmatrix}$$

(and that the occurring matrix satisfies the parameterized Yang-Baxter equation). Verify (1). Find the analogous matrix if we replace $c^{\text{sm}}(\Omega_I)$'s with $[\overline{\Omega}_I]$'s.