

Motivic Chern classes of Schubert cells and applications.

joint works with P. Aluffi, L. Mihalcea, J. Schürmann

Plan: 0) Chern-Schwartz-MacPherson class in homology

1) Motivic Chern class in K-theory

2) Applications in p-adic group representations.

o) Chern-Schwartz-MacPherson class in homology

two functors: $F(X) =$ constructible functions on an alg. variety X/\mathbb{C}
(can be singular).

$$H_*(X) = \text{Homology of } X.$$

Thus (MacPherson).

$\exists!$ natural transformation

$$c_*: F \rightarrow H_*$$

s.t. if X is projective and smooth, $c_*(1_X) = c(TX) \cap [X]$.

$f: X \rightarrow Y$ proper,

$$\begin{array}{ccc} F(X) & \xrightarrow{f_*} & F(Y) \\ c_* \downarrow & \curvearrowright & \downarrow c_* \\ H_*(X) & \xrightarrow{f_*} & H_*(Y) \end{array}$$

Notations:

G s.s. Lie group/ \mathbb{C} , e.g. $SL(n, \mathbb{C})$

B - Borel subgr. $B = \left\{ \begin{pmatrix} * & & \\ & \dots & \\ & & * \end{pmatrix} \right\}$

T - max. torus. $T = \left\{ \begin{pmatrix} * & & \\ & \dots & \\ & & * \end{pmatrix} \right\}$


W - Weyl group $W = S_n$.

$X := G/B$ flag variety, $X = \left\{ F_i = (F_1 \subseteq F_2 \subseteq \dots \subseteq F_n = \mathbb{C}^n) \mid \dim F_i = i \right\}$

$w \in W$, $X(w)^\circ := BwB/B$ Schubert cell.

$X(w) := \overline{X(w)^\circ}$ Schubert variety.

Ex. $G = SL(2, \mathbb{C})$, $X = \mathbb{P}^1$,



$$C_* (\mathbb{1}_{X(\text{id})}) = [X(\text{id})] = [0], \quad C_* (\mathbb{1}_{X(\infty)}) = C_* (\mathbb{1}_{\mathbb{P}^1}) - C_* (\mathbb{1}_{X(\text{id})}) = [1] + [0]$$

Thus: ① (Aluffi-Mihalcea 16) $C_* (\mathbb{1}_{X(w)})$ is generated by the degenerate Hecke operators.

(Aluffi-Mihalcea-Schurman-5, 17)

② $i: X \hookrightarrow T^*X$,

$$C_* (\mathbb{1}_{X(w)}) = \pm i^* [\text{char}(\mathbb{1}_{X(w)})] \Big|_{h=1} = \pm i^* (\text{Maulik-Oblomkov stable envelope}) \Big|_{h=1}$$

equiv. parameter for $G^* \curvearrowright T^*X$.
dilation.

③ $C_* (\mathbb{1}_{X(w)})$ is a positive class in $H^*(X)$. (conjectured by Aluffi-Mihalcea)

1) Motivic Chern class in K-theory

• Definition.

two functors: $\cdot, Y/\mathbb{C}$, $K^0(\text{Var}/Y) := \{ [Z \xrightarrow{f} Y] \} / \{ [Z \xrightarrow{f} Y] = [u \xrightarrow{f} Y] + [Z \wedge u \xrightarrow{f} Y] \}$

$$\cdot K(Y) := K^0(\text{Coh}(Y))$$

$u \subseteq Z$
open.

Thm (Brasselet-Schurmann-Yokura)

$\exists!$ natural transformation

$$\mu_Y: K^0(\text{Var}/-) \rightarrow K(-)[y], \quad \text{st. if } Y \text{ is smooth,}$$

$$\mu_Y([Y \xrightarrow{id} Y]) = \lambda_Y(T^*Y) := \sum_i y^i [\wedge^i T^*Y]. \quad \text{Here } y \text{ is a formal variable.}$$

Remark: \exists equivariant generalizations. (Fehér-Rimányi-Weber, Aluffi-Mihalcea-Schurmann-?)

• Flag variety setting.

$$T \curvearrowright X = G/B \quad K_T(X) := K^0(\text{Coh}_T(X))$$

$$K_T(\text{pt}) = K^0(\text{Coh}_T(\text{pt})) = K^0(\text{Rep}(T)) = \mathbb{Z}[T].$$

$$\text{Let } \text{MG}_Y(X(\omega)^\circ) := \text{MG}_Y([X(\omega)^\circ \hookrightarrow X]) \in K_T(X)[Y].$$


$$\text{Ex. } G = \text{SL}(2, \mathbb{C}), \quad \text{MG}_Y(X(\text{id})^\circ) = [\mathcal{O}_0]$$

$$\text{MG}_Y(X(\omega)^\circ) = \text{MG}_Y(\mathbb{P}^1) - \text{MG}_Y(X(\text{id})^\circ) = \chi_Y(T^*\mathbb{P}^1) - [\mathcal{O}_0].$$

Demazure operators.

e.g. $d_i = \delta_i - \delta_{i+1}$

α_i : simple root, $B \subseteq P_i$ minimal parabolic



$$\pi_i: G/B \rightarrow G/P_i$$

BGG operator $\partial_i \cdot = \pi_i^* \pi_{i*} \in K_T(X)$.

then $\partial_i^2 = \partial_i$, braid relation.

$$\partial_i([\mathcal{O}_{X(w)}]) = [\mathcal{O}_{X(w s_i)}] \quad \text{if } w s_i > w.$$

Demazure-Lusztig operator.

$$\forall \lambda \in X^*(T), \quad L_\lambda := G \times_B G_\lambda$$

$$\downarrow$$

$$X$$

Let $T_i = (1 + y L_{\alpha_i}) \partial_i - \text{id}$, $(T_i + 1)(T_i + y) = 0$, \rightarrow affine Hecke alg.
Braid relations.

Thm: (Aluffi-Mihalcea-Schurmann.-S, 2019).

1). $T_i (MG(X(\omega^\circ))) = MG(X(\omega_{\xi_i}^\circ))$ if $\omega_{\xi_i} > \omega$.

2) $i: X \hookrightarrow T^*X$, $MG(X(\omega^\circ)) = i^* \text{gr}[\mathbb{Q}_{X(\omega^\circ)}^H] = i^* \left(\begin{array}{l} K\text{-theoretic MO} \\ \text{stable envelope} \end{array} \right)$.

constant \uparrow mixed Hodge module.

Fukaya-Rimanyi-Weber

● Smoothness Criterion.

Thm (Kumar). $u \leq w \in W$

$$X(w) \text{ smooth at } u \Leftrightarrow [X(w)]|_u = \prod_{\substack{\alpha > 0 \\ w \leq \alpha \leq w}} (-u\alpha) \in H_T^*(pt).$$

\cap
 $X(w)$

Thm: (Aluffi-Mihalcea-Schurmann.-S, 2019).

$u \leq w \in W$.

$$X(w) \text{ smooth at } u \Leftrightarrow \text{McG}(X(w))|_u = \prod_{\substack{\alpha > 0 \\ u \leq \alpha \leq w}} (1 - e^{u\alpha}) \cdot \prod_{\substack{\alpha > 0 \\ u \leq \alpha \leq w}} (1 + y e^{u\alpha}).$$

(\Rightarrow use property of $\text{McG}(X(w))$,

$\Leftarrow y=0$, $\text{McG}(X(w))|_{y=0} = [\mathcal{O}_{X(w)}(-\partial X(w))]$, take Chern character)

2) Applications in p -adic group representations.

• Bump-Nakassji-Naruse conf.

F non-archimedean local field. $\mathcal{O}_F \subseteq F$, $k_F = \text{residue field} = \overline{\mathbb{F}}_q$, a finite field.

$G^\vee = \text{Langlands dual group} / F$, $T^\vee \subseteq B^\vee \subseteq G^\vee$, $I = \text{Iwahori-subgroup}$, $I \subseteq G^\vee(\mathcal{O}_F)$
 \downarrow \downarrow
 $B^\vee(k_F) \subseteq G^\vee(k_F)$

τ — an unramified char. of T^\vee ($\Leftrightarrow \tau \in T$)

Principal series representation $\text{Ind}_{B^\vee(F)}^{G^\vee(F)}(\tau) \hookrightarrow G^\vee(F)$

Iwahori-Hecke alg.

Let $I(\tau) := \left(\text{Ind}_{B^\vee(F)}^{G^\vee(F)}(\tau) \right)^I \hookrightarrow \mathbb{C}[I \backslash G^\vee(F) / I]$

Two bases in $I(\tau)$: ① $\{\varphi_w \mid w \in W\}$, $\varphi_w = \mathbb{1}_{B^-(F)wI}$.

② $\{f_w \mid w \in W\}$. Casselman basis (defined using the
intertwiners; eigenbasis
for the lattice part of the
Iwahori-Hecke alg.).

define matrix coefficients $m_{u,w}$ by

$$\phi_u := \sum_{w \succ u} \varphi_w = \sum m_{u,w} \cdot f_w$$

Gindikin-Karpelevich formula: $m_{id,w} = \prod_{\substack{\alpha > 0 \\ S \alpha \leq w}} \frac{1 - q^{-\alpha}(\tau)}{1 - e^{\alpha}(\tau)}$.

Let $w_0 \in W$ be the longest element.

Conj (Bump-Nakasuj):

Assume the Dynkin diagram of G is simply-laced.

1). For any $u \leq w \in W$, the Kazhdan-Lusztig poly. $P_{w, w^{-1}, u, u^{-1}}(q) = 1$

$$\Leftrightarrow M_{uw} = \prod_{\alpha > 0} \frac{1 - q^{-1} e^{\alpha}(\tau)}{1 - e^{\alpha}(\tau)}.$$

$u \leq s_{\alpha} w < w$

2). $\prod_{\alpha > 0} (1 - e^{\alpha}) \cdot m_{u, w}$ has no pole on $T(\mathbb{C})$.

$u \leq s_{\alpha} w < w$

Refinement (Naruse). Remove the condition: " G is simply-laced",

The opposite Schubert variety $Y(u) := \overline{B^{-} u B / B} \Leftrightarrow M_{uw} = \prod_{\alpha > 0} \frac{1 - q^{-1} e^{\alpha}(\tau)}{1 - e^{\alpha}(\tau)}$.

is smooth at the point $wB \in Y(u)$.

$u \leq s_{\alpha} w < w$

Thm (Aluffi - Mihalec - Schwarmann - S, 19).

The conjectures hold.

idea: ① G simply laced,

$P_{u,w}(q) = 1 \iff X(w)$ is smooth at $uB \in G/B$.

② $\exists!$ affine Hecke algebra module isomorphism

$$K_T(X) \otimes_{K_T(pt)} \mathbb{C}_T \cong I(\mathbb{C})$$

s.t. $\left\{ \text{dual basis of } \mathcal{M}_G(X(w)^o) \right\} \longleftrightarrow \{ \varphi_w \}$

$$\left\{ \text{fixed point basis} \right\} \longleftrightarrow \{ f_w \}$$

$$\begin{array}{c} T \in T \\ K_T(pt) = \mathbb{Z}[T] \xrightarrow{\text{ev}_T} \mathbb{C}_T \end{array}$$

• Iwahori-Whittaker functions.

σ - an unramified principal character of $N^\vee(\mathbb{F})$ $B^\vee = T^\vee \cdot N^\vee$

Whittaker functional:

$$L: \text{Ind}_{B^\vee(\mathbb{F})}^{G^\vee(\mathbb{F})} \tau \rightarrow \mathbb{C}, \quad \text{s.t.} \quad L(n\phi) = \sigma(n) \cdot L(\phi), \quad n \in N^\vee(\mathbb{F}).$$

For any $f \in \text{Ind}_{B^\vee(\mathbb{F})}^{G^\vee(\mathbb{F})} \tau$,

define $W_\tau(f): G^\vee(\mathbb{F}) \rightarrow \mathbb{C}$

$$g \mapsto L(g \cdot f).$$

Spherical Whittaker function. $W_z \left(\sum_{\omega} \varphi_{\omega} \right) : G^{\vee}(F) \rightarrow \mathbb{C}$.

Thm: (Casselman-Shalika formula.)

μ dominant coweight of G^{\vee} , ϖ -uniformizer of \mathcal{O}_F

$$W_{\tau^{-1}} \left(\sum_{\omega} \varphi_{\omega} \right) (\varpi^{-\mu}) = (*) \prod_{\alpha > 0} (1 - q^{-\alpha} e^{-\alpha}(\varpi)) \cdot \chi_{\mu}(\varpi).$$

\uparrow
char. of ir. highest weight μ
rep. of G .

Remark: $\chi_{\mu} = \chi(G/B^-, G_{\mathbb{A}^{\times}}^{\times} \cdot C_{\mu}) := \sum (-1)^i \text{ch}_T H^i(G/B^-, L_{\mu}) \in K_T(pt)$
 \uparrow equivariant Euler character.
 \parallel
 L_{μ}

Iwahori-Whittaker functions: $W_z(\varphi_w): G^v(F) \rightarrow \mathbb{C}$.

Thy: (Mihalcea - S. 19) μ dominant coweight of G^v ,

$$W_z^{-1}(\varphi_w)(w^{-\mu}) = (*) \prod_{\alpha > 0} (1 + y e^{-\alpha}(z)).$$

$$\chi \left(\frac{G/B^-, L_{\mu} \otimes \frac{M_G([B^- w B^- \hookrightarrow G/B^-])}{\lambda_y(G/B^-)}}{z} \right) (z)$$

Segre-type class.

Remark: 1) Summing over $w \in W$, we get the Casselman-Shalika formula.

2) proof uses work of Bruinaker-Bump-Licata.

3) can also be computed from colored vertex models
(Bruinaker-Buciumas-Bump-Gustafsson).

Thank you!