

# ICERM Lecture 3

---

(geodesic planes in  
 $\infty$ -vol hyp mflds)

Hee Oh

(Yale University) May 17, 2023

---



---

---

$$G = SO^{\circ}(n, 1) = \text{Isom}^+(H^n) \quad n \geq 3$$

$\Gamma < G$  convex cocompact

$$\text{RFM} = \{[g] \in \frac{G}{\Gamma} \mid g^{\pm} \in \Lambda\} \subset \text{RF}_{\Gamma} M = \{[g] \mid g^{\pm} \in \Lambda\} \subset \frac{G}{\Gamma}$$



$$N = \left\{ \begin{pmatrix} 1 & x & * \\ & I & x^t \\ & & 1 \end{pmatrix} \mid x \in \mathbb{R}^{n-1} \right\} \quad \text{max unipotent subgrp } < G$$

$$U \cong \mathbb{R}^{k-1} < N \quad \text{conn.}$$

$$H(U) = \langle U, U^t \rangle \cong SO^0(k, 1)$$

$$\overline{xH(U)} \quad ? \quad \overline{xU} \quad ?$$

$$\mathcal{L}_{H(U)} = \left\{ L = H(\hat{U})C \mid U < \hat{U} < N \right. \\ \left. \cong H(\hat{U})C \text{ closed for some } z \in \mathbb{R}F_+M \right\}$$

↖ any reductive subgrp  $\supset H(U)$

$$\mathcal{L}_U = \left\{ vLv^{-1} \mid L \in \mathcal{L}_U, v \in N \right\}$$

↖ any reductive subgrp  $\supset U$

Thm (MMO  $n=3$ , LO  $n \geq 4$ )

$M = \mathbb{P} \setminus \mathbb{H}^n$  CC hyp mfld with Fuchsian ends

(1)  $H(U)$ -orbit closures

$\forall x \in RFM,$

$$\overline{xH(U)} = xL \cap RF_+M \cdot H(U)$$

for some  $L \in \mathcal{L}_{H(U)}$

(2)  $U$ -orbit closures

$\forall x \in RF_+M,$

$$\overline{xU} = xL \cap RF_+M$$

for some  $L \in \mathcal{L}_U$

(3) Equidistributions

$x_i L_i$  max. closed orbits

$x_i \in RF_+M$   
 $L_i \in \mathcal{L}_U$

$$\lim_{i \rightarrow \infty} x_i L_i \cap RF_+M = RF_+M$$

For  $i = 1, 2, 3,$

$(\bar{i})_m$  holds if  $(i)$  is true for all  
 $U < N$  with  $\text{codim} \leq m$

$m=0$  (1) & (3) trivial

(2) minimality of  $N$ -action

$$(2)_m + (3)_m \Rightarrow (1)_{m+1} + (2)_m + (3)_m$$

$$\Rightarrow (1)_{m+1} + (2)_{m+1} + (3)_m$$

$$\Rightarrow (2)_{m+1} + (3)_{m+1}$$

Rmk If  $\text{vol}(M) < \infty$ ,  $(3)_m$  is not needed  
in the induction pf.

What is special about  $\mathbb{C}$  mflds with Fuchsian ends?

"Recurrence of  $U = \{u_t \mid t \in \mathbb{R}\}$ -orbits"

If  $\frac{G}{\Gamma}$  cpt,  $xU$  remains in a cpt subset

If  $\text{vol}(\frac{G}{\Gamma}) < \infty$ , Dani-Margulis

$\forall x \in \frac{G}{\Gamma}$ ,  $\forall \varepsilon > 0$ ,  $\exists$  cpt  $\mathcal{C}$  s.t.

$$\frac{1}{2T} |\{t \in [-T, T] \mid xu_t \in \mathcal{C}\}| \geq (1-\varepsilon)$$

If  $\text{vol}(\frac{G}{\Gamma}) = \infty$ , for a.e  $x$ ,

$\forall$  cpt  $\mathcal{C} \subset \frac{G}{\Gamma}$ ,

$$\frac{1}{2T} |\{t \in [-T, T] \mid xu_t \in \mathcal{C}\}| \rightarrow 0$$

Prop  $M = \mathbb{R} \setminus \mathbb{H}^n$  Fuchsian ends


$\exists k > 1$  s.t.  $\forall x \in \text{RFM}$ ,

$T(x) = \{t \in \mathbb{R} \mid x U_t \in \text{RFM}\}$  is  $k$ -thick,


i.e.,  $\forall r > 0$ ,

$$T(x) \cap ([-kr, -r] \cup [r, kr]) \neq \emptyset$$

Pf)  $\varepsilon_0 = \inf_{i \neq j} d(\text{hall } B_i, \text{hall } B_j)$   $S^2 - \Lambda = \cup B_i$




$\mathbb{H}^2$



$0, \infty \in \Lambda$

$U = \mathbb{R} = \{\text{real axis}\}$




$\{t \mid [e] U_t \in \text{RFM}\}$

$\parallel$

$\{t \mid U_t^- = t \in \Lambda\}$

$= \mathbb{R} \cap \Lambda$

6



$$d_{\mathbb{H}^2} \left( \text{---} \begin{array}{c} \text{---} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \text{---} \end{array} \right) \geq d_{\mathbb{H}^2}(\text{hull } B_i, \text{hull } B_j) \\ \forall \varepsilon_0$$

$\neq$

## Unipotent blowup lemma

Let  $g_n \rightarrow e$  in  $G - N(U)$

$T \subset \mathbb{R} \cong U$   $K$ -thick subset

$\limsup_{n \rightarrow \infty} T g_n U$  contains  $g \in N(U) - U$ .  
 $d(g, e) < \varepsilon$

Spse  $y U_n \rightarrow x \in \overline{y U}$

$\parallel$   
 $x g_n$

$g_n \notin N(U)$   
 $g_n \rightarrow e$



Since  $T(x) = \{t \in \mathbb{R} = U \mid x u_t \in \text{RFM}\}$   
is  $k$ -thick,

$\exists u_{t_n} \in T(x)$  and  $u_{s_n} \in U$

s.t.  $u_{t_n} g_n u_{s_n} \rightarrow g \in N(U) - U$ .

$$x g_n u_{s_n} = x u_{t_n} (u_{t_n} g_n u_{s_n})$$

$\in \text{RFM}$

$$\rightarrow \exists g \in \overline{yU}$$

This is good enough for  $n=3$

to show  $P$  is closed or dense

in  $\mathbb{H}^3$

In higher  $\dim \geq 4$ , we also need

## Avoidance principle

$\mathcal{Q}(U) = \{x \in \mathbb{R}F_t M \mid xU \text{ is not contained in any closed orbit of } L \neq G\}$

$$\mathcal{J}(U) = \mathbb{R}F_t M - \mathcal{Q}(U)$$

Want: If  $x \in \mathcal{Q}(U)$ ,  $\overline{xU} = \mathbb{R}F_t M$

For this, we need to understand

$$T^*(x) = \{t \in \mathbb{R} \mid xU_t \in \mathbb{R}F_t M, xU_t \notin \mathcal{J}(U)\}$$

# Avoidance Thm (Lee-O.)

$M = \mathbb{P}^1 \setminus \mathbb{H}^n$  Fuchsian ends.

$\exists$  cpt subsets  $E_1 \subset E_2 \subset \dots$  s.t

$$\mathcal{G}(U) \cap \text{RFM} = \bigcup_{i \geq 1} E_i \quad \text{s.t}$$

$\forall x \in \mathcal{G}(U) \cap \text{RFM}, \quad \forall i \geq 1,$

$\exists$  open  $O_i \supset E_i$  s.t

$$T(x) = \{t \in \mathbb{R} \mid \chi_{U_t} \in \text{RFM} - O_i\}$$

is  $k_0$ -thick.

Thank You !!

