

Beyond Algebraic Models of Stringy Spacetime

Tristan Hübsch

@ ICERM: 2026.10.01

Departments of Physics & Astronomy and Mathematics, Howard University, Washington DC

Department of Physics, Faculty of Natural Sciences, Novi Sad University, Serbia

Department of Mathematics, University of Maryland, College Park, MD

Berglund

Algebraic Models of Stringy Spacetime



Tristan Hübsch

@ ICERM: 2026.10.01

w/Per Berglund

🙏: *Mikiya Masuda*

Departments of Physics & Astronomy and Mathematics, Howard University, Washington DC

Department of Physics, Faculty of Natural Sciences, Novi Sad University, Serbia

Department of Mathematics, University of Maryland, College Park, MD

Beyond Algebraic Models (BAM)


Playbill

A Decade Ago
A Family of Scrolls

and Their Offspring
...in the Mirror

- * "It doesn't matter what it's called,
 - * ...as long as it has substance."
- S.-T. Yau






CY n -folds in non-Fano Spaces

How hard can it be?

CY $\subset$ some “nice enough” Ambient space

- Reduce to 0 dimensions: $\mathbb{P}^4[5] \rightarrow \mathbb{P}^3[4] \rightarrow \mathbb{P}^2[3] \rightarrow \mathbb{P}^1[2]$







CY n -folds in non-Fano Spaces

How hard can it be?

CY $\subset$ some “nice enough” Ambient space

- Reduce to 0 dimensions: $\mathbb{P}^4[5] \rightarrow \mathbb{P}^3[4] \rightarrow \mathbb{P}^2[3] \rightarrow \mathbb{P}^1[2]$






CY n -folds in non-Fano Spaces


How hard can it be?

CY $\subset$ some “nice enough” Ambient space





- Reduce to 0 dimensions: $\mathbb{P}^4[5] \rightarrow \mathbb{P}^3[4] \rightarrow \mathbb{P}^2[3] \rightarrow \mathbb{P}^1[2]$




$$\mathcal{X} \in A[c_1]$$



Counting with multiplicity





A Decade Ago...

Classical Constructions


• E.g:
$$X_m \in \left[\begin{array}{c|c|c} \mathbb{P}^4 & 1 & 4 \\ \hline \mathbb{P}^1 & m & 2-m \end{array} \right]_{-168}^{(2,86)}$$

smooth $\mathbb{R}$ models

$$\begin{aligned} b_2 = 2 = h^{1,1} & \text{ dim. space of Kähler classes} \\ \frac{1}{2}b_3 - 1 = 86 = h^{2,1} & \text{ dim. space of cpx structures} \\ -168 = \chi = 2(h^{1,1} - h^{2,1}) & \text{ the Euler \#} \end{aligned}$$

• Zero-set of $p(x, y) = 0$, $\deg[p] = \binom{1}{m}$ & $q(x, y) = 0$, $\deg[q] = \binom{4}{2-m}$

• Generic $\{p=0\} \cap \{q=0\}$ smooth; $\deg_{\mathbb{P}^n}[p] + \deg_{\mathbb{P}^n}[q] = n+1 \Rightarrow c_1 = 0$



A Decade Ago...

Classical Constructions

• E.g: $X_m \in \left[\begin{array}{c|c} \mathbb{P}^4 & 1 \\ \mathbb{P}^1 & m \end{array} \right]_{-168}^{(2,86)}$


smooth $\mathbb{R}$ models

$$\begin{aligned} b_2 = 2 = h^{1,1} & \text{ dim. space of Kähler classes} \\ \frac{1}{2}b_3 - 1 = 86 = h^{2,1} & \text{ dim. space of cpx structures} \\ -168 = \chi = 2(h^{1,1} - h^{2,1}) & \text{ the Euler \#} \end{aligned}$$

• Zero-set of $p(x, y) = 0$, $\deg[p] = \binom{1}{m}$ & $q(x, y) = 0$, $\deg[q] = \binom{4}{2-m}$

• Generic $\{p=0\} \cap \{q=0\}$ smooth; $\deg_{\mathbb{P}^n}[p] + \deg_{\mathbb{P}^n}[q] = n+1 \Rightarrow c_1 = 0$

• Sequentially: $X_m \xrightarrow{q=0} (F_m^{(4)} \xrightarrow{p=0} \mathbb{P}^4 \times \mathbb{P}^1)$



A Decade Ago...

Classical Constructions

• E.g: $X_m \in \left[\begin{array}{c|c} \mathbb{P}^4 & 1 \\ \mathbb{P}^1 & m \end{array} \right]_{-168}^{4, 2-m} (2, 86)$

smooth $\mathbb{R}$ models

special? symplectic

$$\begin{aligned} b_2 = 2 = h^{1,1} & \text{ dim. space of Kähler classes} \\ \frac{1}{2}b_3 - 1 = 86 = h^{2,1} & \text{ dim. space of cpx structures} \\ -168 = \chi = 2(h^{1,1} - h^{2,1}) & \text{ the Euler \#} \end{aligned}$$

• Zero-set of $p(x, y) = 0$, $\deg[p] = \binom{1}{m}$ & $q(x, y) = 0$, $\deg[q] = \binom{4}{2-m}$

• Generic $\{p=0\} \cap \{q=0\}$ smooth; $\deg_{\mathbb{P}^n}[p] + \deg_{\mathbb{P}^n}[q] = n+1 \Rightarrow c_1 = 0$


• Sequentially: $X_m \xrightarrow{q=0} (F_m^{(4)} \xrightarrow{p=0} \mathbb{P}^4 \times \mathbb{P}^1)$

• Chern: $c = \frac{(1+J_1)^5(1+J_2)^2}{(1+J_1+mJ_2)(1+4J_1+(2-m)J_2)}$ for $F_m^{(4)}$ and X_m

• Then: $C_{4-k}^{(c)}[(aJ_1+bJ_2)^k] = \tilde{C}_k(4b+ma) = \text{diffeomorphism invariants}$

• e.g., $\int_{F_m} c_1 c_2 [(aJ_1+bJ_2)] = \int_{X_m} c_2 [(aJ_1+bJ_2)]$, but also $\int_{F_m} c_2 [(aJ_1+bJ_2)^2]$, ...

• So, $F_m^{(4)} \approx_{\mathbb{R}} F_{m \pmod{4}}^{(4)}$ & $X_m \approx_{\mathbb{R}} X_{m \pmod{4}}$: 4 diffeomorphism types



A Decade Ago...

Classical Constructions

• E.g: $X_m \in \left[\begin{array}{c|c} \mathbb{P}^4 & 1 \\ \mathbb{P}^1 & m \end{array} \right]_{-168}^{4, 2-m}^{(2,86)}$

smooth $\mathbb{R}$ models

special? symplectic

$$\begin{aligned} b_2 = 2 = h^{1,1} & \text{ dim. space of Kähler classes} \\ \frac{1}{2}b_3 - 1 = 86 = h^{2,1} & \text{ dim. space of cpx structures} \\ -168 = \chi = 2(h^{1,1} - h^{2,1}) & \text{ the Euler \#} \end{aligned}$$

• Zero-set of $p(x, y) = 0$, $\deg[p] = \binom{1}{m}$ & $q(x, y) = 0$, $\deg[q] = \binom{4}{2-m}$

• Generic $\{p=0\} \cap \{q=0\}$ smooth; $\deg_{\mathbb{P}^n}[p] + \deg_{\mathbb{P}^n}[q] = n+1 \Rightarrow c_1 = 0$

• Sequentially: $X_m \xrightarrow{q=0} (F_m^{(4)} \xrightarrow{p=0} \mathbb{P}^4 \times \mathbb{P}^1)$


• Chern: $c = \frac{(1+J_1)^5(1+J_2)^2}{(1+J_1+mJ_2)(1+4J_1+(2-m)J_2)}$ for $F_m^{(4)}$ and X_m

• Then: $C_{4-k}^{(c)}[(aJ_1+bJ_2)^k] = \tilde{C}_k(4b+ma) = \text{diffeomorphism invariants}$

• e.g., $\int_{F_m} c_1 c_2 [(aJ_1+bJ_2)] = \int_{X_m} c_2 [(aJ_1+bJ_2)]$, but also $\int_{F_m} c_2 [(aJ_1+bJ_2)^2]$, ...

• So, $F_m^{(4)} \approx_{\mathbb{R}} F_{m \pmod{4}}^{(4)}$ & $X_m \approx_{\mathbb{R}} X_{m \pmod{4}}$: 4 diffeomorphism types

• ...so, $m = 0, 1, 2, \dots$



A Decade Ago...

Classical Constructions

• E.g:

$$X_m \in \left[\begin{array}{c|c} \mathbb{P}^4 & 1 \\ \hline \mathbb{P}^1 & m \end{array} \right]_{-168}^{4 \quad 2-m} \quad (2,86)$$

smooth $\mathbb{R}$ models

special? symplectic

$$\begin{aligned} b_2 = 2 = h^{1,1} & \text{ dim. space of Kähler classes} \\ \frac{1}{2}b_3 - 1 = 86 = h^{2,1} & \text{ dim. space of cpx structures} \\ -168 = \chi = 2(h^{1,1} - h^{2,1}) & \text{ the Euler \#} \end{aligned}$$

• Zero-set of $p(x, y) = 0$, $\deg[p] = \binom{1}{m}$ & $q(x, y) = 0$, $\deg[q] = \binom{4}{2-m}$

• Generic $\{p=0\} \cap \{q=0\}$ smooth; $\deg_{\mathbb{P}^n}[p] + \deg_{\mathbb{P}^n}[q] = n+1 \Rightarrow c_1 = 0$

• Sequentially: $X_m \xrightarrow{q=0} (F_m^{(4)} \xrightarrow{p=0} \mathbb{P}^4 \times \mathbb{P}^1)$

• Chern: $c = \frac{(1+J_1)^5(1+J_2)^2}{(1+J_1+mJ_2)(1+4J_1+(2-m)J_2)}$ for $F_m^{(4)}$ and X_m

• Then: $C_{4-k}^{(c)}[(aJ_1+bJ_2)^k] = \tilde{C}_k(\underline{4b+ma}) = \text{diffeomorphism invariants}$

• e.g., $\int_{F_m} c_1 c_2 [(aJ_1+bJ_2)] = \int_{X_m} c_2 [(aJ_1+bJ_2)]$, but also $\int_{F_m} c_2 [(aJ_1+bJ_2)^2]$, ...

• So, $F_m^{(4)} \approx_{\mathbb{R}} F_{m \pmod{4}}^{(4)}$ & $X_m \approx_{\mathbb{R}} X_{m \pmod{4}}$: 4 diffeomorphism types

• ...so, $m=0, 1, 2, \cdot 3 \Rightarrow \deg[q] = \binom{4}{-1} ?!$

$$q(x, y) \stackrel{?}{\sim} \frac{q_0(x)}{y_0} + \frac{q_1(x)}{y_1}$$



A Decade Ago...

Why Haven't We Thought of This Before?

- $\deg[q] = \binom{4}{-1}$ holomorphic sections?!


- Not *everywhere* on $\mathbb{P}^4 \times \mathbb{P}^1$ — (simple poles)

- but yes on $F_3^{(4)} \subset \mathbb{P}^4 \times \mathbb{P}^1$ — ≥ 105 of 'em!

$$X_m \in \left[\begin{array}{c|c} \mathbb{P}^4 & 1 \\ \mathbb{P}^1 & m \end{array} \right]_{2-m}^{(2,86)} \quad \text{for } m=3$$




[AAGGL:1507.03235 + BH:1606.07420]



A Decade Ago...

Why Haven't We Thought of This Before?



[AAGGL:1507.03235 + BH:1606.07420]

- $\deg[q] = \binom{4}{-1}$ holomorphic sections?!

- Not *everywhere* on $\mathbb{P}^4 \times \mathbb{P}^1$ — (simple poles)


- but yes on $F_3^{(4)} \subset \mathbb{P}^4 \times \mathbb{P}^1$ — ≥ 105 of 'em!

$$X_m \in \left[\begin{array}{c|c} \mathbb{P}^4 & 1 \\ \mathbb{P}^1 & m \end{array} \right]_{2-m}^{(2,86)} \quad \text{for } m=3$$

- How? On $F_3^{(4)}$, $q(x, y) \simeq q(x, y) + \lambda \cdot p(x, y) \leftarrow$ equivalence class!

A Decade Ago...

Why Haven't We Thought of This Before?



[AAGGL:1507.03235 + BH:1606.07420]
[+ GvG:1708.00517]

- $\deg[q] = \binom{4}{-1}$ holomorphic sections?!

- Not *everywhere* on $\mathbb{P}^4 \times \mathbb{P}^1$ — (simple poles)

- but yes on $F_3^{(4)} \subset \mathbb{P}^4 \times \mathbb{P}^1$ — ≥ 105 of 'em!

$$X_m \in \left[\begin{array}{c|c} \mathbb{P}^4 & 1 \\ \mathbb{P}^1 & m \end{array} \right]_{2-m}^{(2,86)} \quad \text{for } m=3$$

Two more choices,

$$\left(\frac{x_0 y_0^2}{y_1^3} - \frac{x_1}{y_0} \right), \left(\frac{x_0}{y_1} - \frac{x_1 y_1^2}{y_0^3} \right)$$

$\Rightarrow 3 \times 35 = 105$ sections

- How? On $F_3^{(4)}$, $q(x, y) \simeq q(x, y) + \lambda \cdot p(x, y) \leftarrow$ equivalence class!

- [Hirzebruch, 1951] $\Rightarrow p = x_0 y_0^3 + x_1 y_1^3$

& $q = c(x) \left[\left(\frac{x_0 y_0}{y_1^2} - \frac{x_1 y_1}{y_0^2} \right) \bmod p_0(x, y) \right] \quad \deg[c] = \binom{3}{0}$

- So, $q_0 = \left[q(x, y) + \frac{\lambda c(x)}{(y_0 y_1)^2} p(x, y) \right]_{\lambda \rightarrow -1} = c(x) \left(-2 \frac{x_1 y_1}{y_0^2} \right)$ in the $y_0 \neq 0$ chart

- & $q_1 = \left[q(x, y) + \frac{\lambda c(x)}{(y_0 y_1)^2} p(x, y) \right]_{\lambda \rightarrow +1} = c(x) \left(2 \frac{x_0 y_0}{y_1^2} \right)$ in the $y_1 \neq 0$ chart

- & $q_1(x, y) - q_0(x, y) = 2 \frac{c(x)}{(y_0 y_1)^2} p(x, y) = 0$ on $F_3 := \{p(x, y) = 0\}$


- [GvG, 1708.00517] *scheme-th.* “generalized complete intersections”

Reverse-engineered: Mayer-Vietoris sequence & “patching” of the two charts

Wu-Yang
magnetic
monopole

A Decade Ago...

Why Haven't We Thought of This Before?



[AAGGL:1507.03235 + BH:1606.07420]
[+ GvG:1708.00517]

- $\deg[q] = \binom{4}{-1}$ holomorphic sections?!

- Not *everywhere* on $\mathbb{P}^4 \times \mathbb{P}^1$ — (simple poles)

- but yes on $F_3^{(4)} \subset \mathbb{P}^4 \times \mathbb{P}^1$ — ≥ 105 of 'em!

$$X_m \in \left[\begin{array}{c|c} \mathbb{P}^4 & 1 \\ \hline \mathbb{P}^1 & m \end{array} \begin{array}{c} 4 \\ 2-m \end{array} \right]_{-168}^{(2,86)} \text{ for } m=3$$

Two more choices,

$$\left(\frac{x_0 y_0^2}{y_1^3} - \frac{x_1}{y_0} \right), \left(\frac{x_0}{y_1} - \frac{x_1 y_1^2}{y_0^3} \right)$$

- How? On $F_3^{(4)}$, $q(x, y) \simeq q(x, y) + \lambda \cdot p(x, y) \leftarrow$ equivalence class!

$\Rightarrow 3 \times 35 = 105$ sections

- [Hirzebruch, 1951] $\Rightarrow p = x_0 y_0^3 + x_1 y_1^3$

& $q = c(x) \left[\left(\frac{x_0 y_0}{y_1^2} - \frac{x_1 y_1}{y_0^2} \right) \bmod p_0(x, y) \right] \deg[c] = \binom{3}{0}$

- So, $q_0 = \left[q(x, y) + \frac{\lambda c(x)}{(y_0 y_1)^2} p(x, y) \right]_{\lambda \rightarrow -1} = c(x) \left(-2 \frac{x_1 y_1}{y_0^2} \right)$ in the $y_0 \neq 0$ chart

- & $q_1 = \left[q(x, y) + \frac{\lambda c(x)}{(y_0 y_1)^2} p(x, y) \right]_{\lambda \rightarrow +1} = c(x) \left(2 \frac{x_0 y_0}{y_1^2} \right)$ in the $y_1 \neq 0$ chart

- & $q_1(x, y) - q_0(x, y) = 2 \frac{c(x)}{(y_0 y_1)^2} p(x, y) = 0$ on $F_3 := \{p(x, y) = 0\}$

- [GvG, 1708.00517] *scheme-th.* “generalized complete intersections”

Reverse-engineered: Mayer-Vietoris sequence & “patching” of the two charts

2026: Borisov: no schemes needed—rank-2 v.bdl & codim-2 0-locus

Wu-Yang
magnetic
monopole

A Family of Scrolls


...in well-tempered counterpoint

[BH:1606.07420, 1611.10300 & 2205.12827]

+more

• For $\left\{ 0 \stackrel{!}{=} \underbrace{x_0 y_0^m + x_1 y_1^m}_{:= p_0(x, y)} + \sum_{a=2}^n \sum_{\ell=1}^{m-1} \epsilon_{a,\ell} x_a y_0^{m-\ell} y_1^\ell \right\} = F_{m;\epsilon}^{(n)} \in \left[\begin{array}{c} \mathbb{P}^n \\ \mathbb{P}^1 \end{array} \middle\| \begin{array}{c} 1 \\ m \end{array} \right]$


even $p_0(x, y)$ is transverse, $p_0^{-1}(0)$ is smooth



A Family of Scrolls

...in well-tempered counterpoint

$p_0^{-1}(0) \cap \mathfrak{z}^{-1}(0)$ is smooth
 $dp_0(x, y) \wedge d\mathfrak{z}(x, y) \neq 0$



[BH:1606.07420, 1611.10300 & 2205.12827]

+more

For $\left\{ 0 \stackrel{!}{=} \underbrace{x_0 y_0^m + x_1 y_1^m}_{:= p_0(x, y)} + \sum_{a=2}^n \sum_{\ell=1}^{m-1} \epsilon_{a,\ell} x_a y_0^{m-\ell} y_1^\ell \right\} = F_{m;\epsilon}^{(n)} \in \left[\begin{array}{c} \mathbb{P}^n \\ \mathbb{P}^1 \end{array} \middle| \begin{array}{c} 1 \\ m \end{array} \right]$

even $p_0(x, y)$ is transverse, $p_0^{-1}(0)$ is smooth

The central ($\epsilon = 0$) member of the family is a Hirzebruch scroll F_m :

Directrix: $\mathfrak{z}(x, y) := \left[\left(\frac{x_0}{y_1^m} - \frac{x_1}{y_0^m} \right) + \frac{\lambda}{(y_0 y_1)^m} p_0(x, y) \right]$ degree- $\left(-\frac{1}{m}\right)$

On $F_m^{(n)}$: $p_0(x, y) = x_0 y_0^m + x_1 y_1^m = 0 \Rightarrow x_0 = -x_1 (y_1 / y_0)^m$, @ $y_0 \neq 0$

So, $x_1 \rightarrow X_1 = \mathfrak{z}(x, y)$ & $(X_i, i=2, \dots) = (x_2, \dots, x_n; y_0, y_1)$

Key: $\det \left[\frac{\partial(p_0(x, y), \mathfrak{z}(x, y), x_2, \dots; y_0, y_1)}{\partial(x_0, x_1, x_2, \dots; y_0, y_1)} \right] = \text{const.}$


$\mathbb{P}^4 \times \mathbb{P}^1$ bi-degree $\rightarrow$ toric $(\mathbb{C}^\times)^2$ -action:

X_1	X_2	X_3	X_4	X_5	X_6	
1	1	1	1	0	0	$\leftarrow \mathbb{P}^4$
$-m$	0	0	0	1	1	$\leftarrow \mathbb{P}^1$

A Family of Scrolls

...in well-tempered counterpoint

$p_0^{-1}(0) \cap \mathfrak{z}^{-1}(0)$ is smooth
 $dp_0(x, y) \wedge d\mathfrak{z}(x, y) \neq 0$



[BH:1606.07420, 1611.10300 & 2205.12827]

+more

For $\left\{ 0 \stackrel{!}{=} \underbrace{x_0 y_0^m + x_1 y_1^m}_{:= p_0(x, y)} + \sum_{a=2}^n \sum_{\ell=1}^{m-1} \epsilon_{a,\ell} x_a y_0^{m-\ell} y_1^\ell \right\} = F_{m;\epsilon}^{(n)} \in \left[\begin{array}{c} \mathbb{P}^n \\ \mathbb{P}^1 \end{array} \middle| \begin{array}{c} 1 \\ m \end{array} \right]$

even $p_0(x, y)$ is transverse, $p_0^{-1}(0)$ is smooth

The central ($\epsilon = 0$) member of the family is a Hirzebruch scroll F_m :

Directrix: $\mathfrak{z}(x, y) := \left[\left(\frac{x_0}{y_1^m} - \frac{x_1}{y_0^m} \right) + \frac{\lambda}{(y_0 y_1)^m} p_0(x, y) \right]$ degree- $\left(-\frac{1}{m}\right)$


On $F_m^{(n)}$: $p_0(x, y) = x_0 y_0^m + x_1 y_1^m = 0 \Rightarrow x_0 = -x_1 (y_1 / y_0)^m$, @ $y_0 \neq 0$

So, $x_1 \rightarrow X_1 = \mathfrak{z}(x, y)$ & $(X_i, i=2, \dots) = (x_2, \dots, x_n; y_0, y_1)$

Key: $\det \left[\frac{\partial(p_0(x, y), \mathfrak{z}(x, y), x_2, \dots; y_0, y_1)}{\partial(x_0, x_1, x_2, \dots; y_0, y_1)} \right] = \text{const.}$

$\mathbb{P}^4 \times \mathbb{P}^1$ bi-degree $\rightarrow$ toric $(\mathbb{C}^\times)^2$ -action:

What about the rest of the $\epsilon_{a,\ell}$ -deformation family?
 (wherein smooth models are all diffeomorphic to each other)



*Reverse-Engineered (Toric) Model

A Family of Scrolls

...with a meandering motif


[BH:1606.07420, 1611.10300 & 2205.12827]

+more

Construction 2.1 Given a degree- $\binom{1}{m}$ hypersurface $\{p_{\vec{\epsilon}}(x, y) = 0\} \subset \mathbb{P}^n \times \mathbb{P}^1$ as in (2.2), construct


$$\deg = \binom{1}{m-r_0-r_1} : \quad \mathfrak{s}_{\vec{\epsilon}}(x, y; \lambda) := \text{Flip}_{y_0} \left[\frac{1}{y_0^{r_0} y_1^{r_1}} p_{\vec{\epsilon}}(x, y) \right] \pmod{p_{\vec{\epsilon}}(x, y)}, \quad \left[\begin{array}{c} \mathbb{P}^n \\ \mathbb{P}^1 \end{array} \middle| \begin{array}{c} 1 \\ m \end{array} \right]$$

progressively decreasing $r_0 + r_1 = 2m, 2m-1, \dots$, and keeping only Laurent polynomials containing both y_0 - and y_1 -denominators but no y_0, y_1 -mixed ones. The “Flip $_{y_i}$ ” operator changes the relative sign of the rational monomials with y_i -denominators. For algebraically independent such sections, restrict to a subset with maximally negative degrees that are not overall (y_0, y_1) -multiples of each other.



A Family of Scrolls

...with a meandering motif



[BH:1606.07420, 1611.10300 & 2205.12827]

+more

Construction 2.1 Given a degree- $\binom{1}{m}$ hypersurface $\{p_{\vec{\epsilon}}(x, y) = 0\} \subset \mathbb{P}^n \times \mathbb{P}^1$ as in (2.2), construct

$$\textcircled{1} \quad \deg = \binom{1}{m-r_0-r_1} : \quad \mathfrak{s}_{\vec{\epsilon}}(x, y; \lambda) := \text{Flip}_{y_0} \left[\frac{1}{y_0^{r_0} y_1^{r_1}} p_{\vec{\epsilon}}(x, y) \right] \pmod{p_{\vec{\epsilon}}(x, y)}, \quad \left[\begin{array}{c|c} \mathbb{P}^n & 1 \\ \hline \mathbb{P}^1 & m \end{array} \right]$$


progressively decreasing $r_0 + r_1 = 2m, 2m-1, \dots$, and keeping only Laurent polynomials containing both y_0 - and y_1 -denominators but no y_0, y_1 -mixed ones. The “Flip $_{y_i}$ ” operator changes the relative sign of the rational monomials with y_i -denominators. For algebraically independent such sections, restrict to a subset with maximally negative degrees that are not overall (y_0, y_1) -multiples of each other.

E.g.: $m=2$
 $p_0 = x_0 y_0^2 + x_1 y_1^2$; $\text{ep}[\alpha] := \text{Table} \left[\frac{1}{y_0^{\alpha-i} y_1^i}, \{i, 0, \alpha\} \right]$; $\text{Expand} /@ (p_0 \{ \text{ep}[5], \text{ep}[4], \text{ep}[3] \})$

$$\textcircled{1} \quad \left\{ \left\{ \frac{x_0}{y_0^3} + \frac{x_1 y_1^2}{y_0^5}, \frac{x_0}{y_0^2 y_1} + \frac{x_1 y_1}{y_0^4}, \frac{x_1}{y_0^3} + \frac{x_0}{y_0 y_1^2}, \frac{x_0}{y_1^3} + \frac{x_1}{y_0^2 y_1}, \frac{x_0 y_0}{y_1^4} + \frac{x_1}{y_0 y_1^2}, \frac{x_0 y_0^2}{y_1^5} + \frac{x_1}{y_1^3} \right\}, \right. \\ \left. \left\{ \frac{x_0}{y_0^2} + \frac{x_1 y_1^2}{y_0^4}, \frac{x_0}{y_0 y_1} + \frac{x_1 y_1}{y_0^3}, \frac{x_1}{y_0^2} + \frac{x_0}{y_1^2}, \frac{x_0 y_0}{y_1^3} + \frac{x_1}{y_0 y_1}, \frac{x_0 y_0^2}{y_1^4} + \frac{x_1}{y_1^2} \right\}, \left\{ \frac{x_0}{y_0} + \frac{x_1 y_1^2}{y_0^3}, \frac{x_0}{y_1} + \frac{x_1 y_1}{y_0^2}, \frac{x_1}{y_0} + \frac{x_0 y_0}{y_1^2}, \frac{x_0 y_0^2}{y_1^3} + \frac{x_1}{y_1} \right\} \right\}$$

A Family of Scrolls

...with a meandering motif



[BH:1606.07420, 1611.10300 & 2205.12827]

+more

Construction 2.1 Given a degree- $\binom{1}{m}$ hypersurface $\{p_{\vec{\epsilon}}(x, y) = 0\} \subset \mathbb{P}^n \times \mathbb{P}^1$ as in (2.2), construct

$$\textcircled{1} \quad \deg = \binom{1}{m-r_0-r_1} : \quad \mathfrak{s}_{\vec{\epsilon}}(x, y; \lambda) := \text{Flip}_{y_0} \left[\frac{1}{y_0^{r_0} y_1^{r_1}} p_{\vec{\epsilon}}(x, y) \right] \pmod{p_{\vec{\epsilon}}(x, y)}, \quad \left[\begin{array}{c} \mathbb{P}^n \\ \mathbb{P}^1 \end{array} \middle| \begin{array}{c} 1 \\ m \end{array} \right]$$

progressively decreasing $r_0 + r_1 = 2m, 2m-1, \dots$, and keeping only Laurent polynomials con-

$\textcircled{2}$ taining both y_0 - and y_1 -denominators but no y_0, y_1 -mixed ones. The “Flip $_{y_i}$ ” operator changes the relative sign of the rational monomials with y_i -denominators. For algebraically independent such sections, restrict to a subset with maximally negative degrees that are not overall (y_0, y_1) -multiples of each other.


E.g.: $m=2$
 $p_0 = x_0 y_0^2 + x_1 y_1^2$; $\text{ep}[\alpha] := \text{Table} \left[\frac{1}{y_0^{\alpha-i} y_1^i}, \{i, 0, \alpha\} \right]$; $\text{Expand} /@ (p_0 \{ \text{ep}[5], \text{ep}[4], \text{ep}[3] \})$

$$\textcircled{1} \quad \left\{ \left\{ \frac{x_0}{y_0^3} + \frac{x_1 y_1^2}{y_0^5}, \frac{x_0}{y_0^2 y_1} + \frac{x_1 y_1}{y_0^4}, \frac{x_1}{y_0^3} + \frac{x_0}{y_0 y_1^2}, \frac{x_0}{y_1^3} + \frac{x_1}{y_0^2 y_1}, \frac{x_0 y_0}{y_1^4} + \frac{x_1}{y_0 y_1^2}, \frac{x_0 y_0^2}{y_1^5} + \frac{x_1}{y_1^3} \right\}, \right.$$

$$\textcircled{2} \quad \left\{ \frac{x_0}{y_0^2} + \frac{x_1 y_1^2}{y_0^4}, \frac{x_0}{y_0 y_1} + \frac{x_1 y_1}{y_0^3}, \frac{x_1}{y_0^2} + \frac{x_0}{y_1^2}, \frac{x_0 y_0}{y_1^3} + \frac{x_1}{y_0 y_1}, \frac{x_0 y_0^2}{y_1^4} + \frac{x_1}{y_1^2} \right\}, \left\{ \frac{x_0}{y_0} + \frac{x_1 y_1^2}{y_0^3}, \frac{x_0}{y_1} + \frac{x_1 y_1}{y_0^2}, \frac{x_1}{y_0} + \frac{x_0 y_0}{y_1^2}, \frac{x_0 y_0^2}{y_1^3} + \frac{x_1}{y_1} \right\} \}$$

A Family of Scrolls

...with a meandering motif



[BH:1606.07420, 1611.10300 & 2205.12827]

+more

Construction 2.1 Given a degree- $\binom{1}{m}$ hypersurface $\{p_{\vec{\epsilon}}(x, y) = 0\} \subset \mathbb{P}^n \times \mathbb{P}^1$ as in (2.2), construct

$$\textcircled{1} \quad \deg = \binom{1}{m-r_0-r_1} : \quad \mathfrak{s}_{\vec{\epsilon}}(x, y; \lambda) := \text{Flip}_{y_0} \left[\frac{1}{y_0^{r_0} y_1^{r_1}} p_{\vec{\epsilon}}(x, y) \right] \pmod{p_{\vec{\epsilon}}(x, y)}, \quad \left[\begin{array}{c|c} \mathbb{P}^n & 1 \\ \hline \mathbb{P}^1 & m \end{array} \right]$$

progressively decreasing $r_0 + r_1 = 2m, 2m-1, \dots$, and keeping only Laurent polynomials con-

$\textcircled{2}$ taining both y_0 - and y_1 -denominators but no y_0, y_1 -mixed ones. The “Flip $_{y_i}$ ” operator changes the relative sign of the rational monomials with y_i -denominators. For algebraically independent such sections, restrict to a subset with maximally negative degrees that are not overall (y_0, y_1) -multiples of each other.


E.g.: $m=2$
 $p_0 = x_0 y_0^2 + x_1 y_1^2$; $\text{ep}[\alpha] := \text{Table} \left[\frac{1}{y_0^{\alpha-i} y_1^i}, \{i, 0, \alpha\} \right]$; $\text{Expand} /@ (p_0 \{ \text{ep}[5], \text{ep}[4], \text{ep}[3] \})$

$$\textcircled{1} \quad \left\{ \left\{ \frac{x_0}{y_0^3} + \frac{x_1 y_1^2}{y_0^5}, \frac{x_0}{y_0^2 y_1} + \frac{x_1 y_1}{y_0^4}, \frac{x_1}{y_0^3} + \frac{x_0}{y_0 y_1^2}, \frac{x_0}{y_1^3} + \frac{x_1}{y_0 y_1^2}, \frac{x_0 y_0}{y_1^4} + \frac{x_1}{y_0 y_1^2}, \frac{x_0 y_0^2}{y_1^5} + \frac{x_1}{y_1^3} \right\}, \right.$$

$$\textcircled{2} \quad \left\{ \frac{x_0}{y_0^2} + \frac{x_1 y_1^2}{y_0^4}, \frac{x_0}{y_0 y_1} + \frac{x_1 y_1}{y_0^3}, \frac{x_1}{y_0^2} + \frac{x_0}{y_1^2}, \frac{x_0 y_0}{y_1^3} + \frac{x_1}{y_0 y_1^2}, \frac{x_0 y_0^2}{y_1^4} + \frac{x_1}{y_1^2} \right\}, \left\{ \frac{x_0}{y_0} + \frac{x_1 y_1^2}{y_0^3}, \frac{x_0}{y_1} + \frac{x_1 y_1}{y_0^2}, \frac{x_1}{y_0} + \frac{x_0 y_0}{y_1^2}, \frac{x_0 y_0^2}{y_1^3} + \frac{x_1}{y_1} \right\} \}$$

A Family of Scrolls

...with a meandering motif



[BH:1606.07420, 1611.10300 & 2205.12827]

+more

Construction 2.1 Given a degree- $\binom{1}{m}$ hypersurface $\{p_{\vec{\epsilon}}(x, y) = 0\} \subset \mathbb{P}^n \times \mathbb{P}^1$ as in (2.2), construct

$$\textcircled{1} \quad \deg = \binom{1}{m-r_0-r_1} : \quad \mathfrak{s}_{\vec{\epsilon}}(x, y; \lambda) := \text{Flip}_{y_0} \left[\frac{1}{y_0^{r_0} y_1^{r_1}} p_{\vec{\epsilon}}(x, y) \right] \pmod{p_{\vec{\epsilon}}(x, y)}, \quad \left[\begin{array}{c} \mathbb{P}^n \\ \mathbb{P}^1 \end{array} \middle| \begin{array}{c} 1 \\ m \end{array} \right]$$

progressively decreasing $r_0 + r_1 = 2m, 2m-1, \dots$, and keeping only Laurent polynomials con-

$\textcircled{2}$ taining both y_0 - and y_1 -denominators but no y_0, y_1 -mixed ones. The “Flip $_{y_i}$ ” operator changes the relative sign of the rational monomials with y_i -denominators. For algebraically independent such sections, restrict to a subset with maximally negative degrees that are not overall (y_0, y_1) -multiples of each other.


E.g.: $m=2$
 $p_0 = x_0 y_0^2 + x_1 y_1^2$; $\text{ep}[\alpha] := \text{Table} \left[\frac{1}{y_0^{\alpha-i} y_1^i}, \{i, 0, \alpha\} \right]$; $\text{Expand} /@ (p_0 \{ \text{ep}[5], \text{ep}[4], \text{ep}[3] \})$

$$\textcircled{1} \quad \left\{ \left\{ \frac{x_0}{y_0^3} + \frac{x_1 y_1^2}{y_0^5}, \frac{x_0}{y_0^2 y_1} + \frac{x_1 y_1}{y_0^4}, \frac{x_1}{y_0^3} + \frac{x_0}{y_0 y_1^2}, \frac{x_0}{y_1^3} + \frac{x_1}{y_0^2 y_1}, \frac{x_0 y_0}{y_1^4} + \frac{x_1}{y_0 y_1^2}, \frac{x_0 y_0^2}{y_1^5} + \frac{x_1}{y_1^3} \right\}, \right.$$

$$\textcircled{2} \quad \left\{ \frac{x_0}{y_0^2} + \frac{x_1 y_1^2}{y_0^4}, \frac{x_0}{y_0 y_1} + \frac{x_1 y_1}{y_0^3}, \frac{x_1}{y_0^2} + \frac{x_0}{y_1^2}, \frac{x_0 y_0}{y_1^3} + \frac{x_1}{y_0 y_1}, \frac{x_0 y_0^2}{y_1^4} + \frac{x_1}{y_1^2}, \left\{ \frac{x_0}{y_0} + \frac{x_1 y_1^2}{y_0^3}, \frac{x_0}{y_1} + \frac{x_1 y_1}{y_0^2}, \frac{x_1}{y_0} + \frac{x_0 y_0}{y_1^2}, \frac{x_0 y_0^2}{y_1^3} + \frac{x_1}{y_1} \right\} \right\}$$

A Family of Scrolls

...with a meandering motif



[BH:1606.07420, 1611.10300 & 2205.12827]

+more

Construction 2.1 Given a degree- $\binom{1}{m}$ hypersurface $\{p_{\vec{e}}(x, y) = 0\} \subset \mathbb{P}^n \times \mathbb{P}^1$ as in (2.2), construct

$$\textcircled{1} \quad \deg = \binom{1}{m-r_0-r_1} : \quad \mathfrak{s}_{\vec{e}}(x, y; \lambda) := \text{Flip}_{y_0} \left[\frac{1}{y_0^{r_0} y_1^{r_1}} p_{\vec{e}}(x, y) \right] \pmod{p_{\vec{e}}(x, y)}, \quad \left[\begin{array}{c|c} \mathbb{P}^n & 1 \\ \hline \mathbb{P}^1 & m \end{array} \right]$$

progressively decreasing $r_0 + r_1 = 2m, 2m-1, \dots$, and keeping only Laurent polynomials con-

$\textcircled{2}$ taining both y_0 - and y_1 -denominators but no y_0, y_1 -mixed ones. The “Flip $_{y_i}$ ” operator changes the relative sign of the rational monomials with y_i -denominators. For algebraically indepen-

$\textcircled{3}$ dent such sections, restrict to a subset with maximally negative degrees that are not overall (y_0, y_1) -multiples of each other.

E.g.: $m=2$
 $p_0 = x_0 y_0^2 + x_1 y_1^2$; $\text{ep}[\alpha] := \text{Table} \left[\frac{1}{y_0^{\alpha-i} y_1^i}, \{i, 0, \alpha\} \right]$; $\text{Expand} /@ (p_0 \{ \text{ep}[5], \text{ep}[4], \text{ep}[3] \})$

$$\textcircled{1} \quad \left\{ \left\{ \frac{x_0}{y_0^3} + \frac{x_1 y_1^2}{y_0^5}, \frac{x_0}{y_0^2 y_1} + \frac{x_1 y_1}{y_0^4}, \frac{x_1}{y_0^3} + \frac{x_0}{y_0 y_1^2}, \frac{x_0}{y_1^3} + \frac{x_1}{x_0 y_1^2}, \frac{x_0 y_0}{y_1^4} + \frac{x_1}{y_0 y_1^2}, \frac{x_0 y_0^2}{y_1^5} + \frac{x_1}{y_1^3} \right\}, \cdot y_1, \cdot y_0 \right\}$$


$$\textcircled{2} \quad \left\{ \frac{x_0}{y_0^2} + \frac{x_1 y_1^2}{y_0^4}, \frac{x_0}{y_0 y_1} + \frac{x_1 y_1}{y_0^3}, \frac{x_1}{y_0^2} + \frac{x_0}{y_1^2}, \frac{x_0 y_0}{y_1^3} + \frac{x_1}{y_0 y_1^2}, \frac{x_0 y_0^2}{y_1^4} + \frac{x_1}{y_1^2}, \left\{ \frac{x_0}{y_0} + \frac{x_1 y_1^2}{y_0^3}, \frac{x_0}{y_1} + \frac{x_1 y_1}{y_0^2}, \frac{x_1}{y_0} + \frac{x_0 y_0}{y_1^2}, \frac{x_0 y_0^2}{y_1^3} + \frac{x_1}{y_1} \right\} \right\}$$

$$\textcircled{3}$$

A Family of Scrolls

...with a meandering motif

$p_0^{-1}(0) \cap \mathfrak{z}^{-1}(0)$ is smooth
 $dp_0(x, y) \wedge d\mathfrak{z}(x, y) \neq 0$



[BH:1606.07420, 1611.10300 & 2205.12827]

+more

Construction 2.1 Given a degree- $\binom{1}{m}$ hypersurface $\{p_{\vec{e}}(x, y) = 0\} \subset \mathbb{P}^n \times \mathbb{P}^1$ as in (2.2), construct

$$\textcircled{1} \quad \deg = \binom{1}{m-r_0-r_1} : \quad \mathfrak{s}_{\vec{e}}(x, y; \lambda) := \text{Flip}_{y_0} \left[\frac{1}{y_0^{r_0} y_1^{r_1}} p_{\vec{e}}(x, y) \right] \pmod{p_{\vec{e}}(x, y)}, \quad \left[\begin{array}{c|c} \mathbb{P}^n & 1 \\ \hline \mathbb{P}^1 & m \end{array} \right]$$

progressively decreasing $r_0 + r_1 = 2m, 2m-1, \dots$, and keeping only Laurent polynomials con-

$\textcircled{2}$ taining both y_0 - and y_1 -denominators but no y_0, y_1 -mixed ones. The “Flip $_{y_i}$ ” operator changes the relative sign of the rational monomials with y_i -denominators. For algebraically indepen-

$\textcircled{3}$ dent such sections, restrict to a subset with maximally negative degrees that are not overall (y_0, y_1) -multiples of each other.

Use this for identification

E.g.: $m=2$
 $p_0 = x_0 y_0^2 + x_1 y_1^2$; $\text{ep}[\alpha] := \text{Table} \left[\frac{1}{y_0^{\alpha-i} y_1^i}, \{i, 0, \alpha\} \right]$; $\text{Expand} /@ (p_0 \{ \text{ep}[5], \text{ep}[4], \text{ep}[3] \})$

$$\textcircled{1} \quad \left\{ \left\{ \frac{x_0}{y_0^3} + \frac{x_1 y_1^2}{y_0^5}, \frac{x_0}{y_0^2 y_1} + \frac{x_1 y_1}{y_0^4}, \frac{x_1}{y_0^3} + \frac{x_0}{y_0 y_1^2}, \frac{x_0}{y_1^3} + \frac{x_1}{y_0^2 y_1^2}, \frac{x_0 y_0}{y_1^4} + \frac{x_1}{y_0 y_1^2}, \frac{x_0 y_0^2}{y_1^5} + \frac{x_1}{y_1^3} \right\}, \cdot y_1, \cdot y_0 \right\}$$

$$\textcircled{2} \quad \left\{ \frac{x_0}{y_0^2} + \frac{x_1 y_1^2}{y_0^4}, \frac{x_0}{y_0 y_1} + \frac{x_1 y_1}{y_0^3}, \frac{x_1}{y_0^2} + \frac{x_0}{y_1^2}, \frac{x_0 y_0}{y_1^3} + \frac{x_1}{y_0 y_1^2}, \frac{x_0 y_0^2}{y_1^4} + \frac{x_1}{y_1^2}, \left\{ \frac{x_0}{y_0} + \frac{x_1 y_1^2}{y_0^3}, \frac{x_0}{y_1} + \frac{x_1 y_1}{y_0^2}, \frac{x_1}{y_0} + \frac{x_0 y_0}{y_1^2}, \frac{x_0 y_0^2}{y_1^3} + \frac{x_1}{y_1} \right\} \right\}$$

finds $\mathfrak{z}(x, y) = \left(\frac{x_0}{y_1^2} - \frac{x_1}{y_0^2} \right) \pmod{(x_0 y_0^2 + x_1 y_1^2)}$; $\deg = \binom{1}{-2}$, $[\mathfrak{z}^{-1}(0)] = [J_1] - 2[J_2]$.

THE exceptional section

THE hallmark
 exceptional divisor
 $[S]^2 = -1$ of $F_2^{(2)}$

A Family of Scrolls

...with cousins joining in...

• Deform: $p_1(x, y) = x_0 y_0^5 + x_1 y_1^5 + x_2 y_0 y_1^4$



• Now: $\mathfrak{s}_{1,1}(x, y) = \frac{x_0 y_0}{y_1^5} + \frac{x_2}{y_1^4} - \frac{x_1}{y_1^4}$ & $\mathfrak{s}_{1,2}(x, y) = \frac{x_0}{y_1} - \frac{x_2}{y_0} - \frac{x_1 y_1^4}{y_0^5}$

• & $\det \left[\frac{\partial(p_1, \mathfrak{s}_{1,1}, \mathfrak{s}_{1,2}, x_3, \dots; y_0, y_1)}{\partial(x_0, x_1, x_2, x_3, \dots; y_0, y_1)} \right] = \text{const.}$

toric $F_{(4,1,0,\dots)}^{(n)}$


X_1	X_2	X_3	X_4	X_5	X_6	
1	1	1	1	0	0	$\leftarrow \mathbb{P}^4$
-4	-1	0	0	1	1	$\leftarrow \mathbb{P}^1$

the central
($\epsilon_{a,\ell}=0$)
model



[BH:1606.07420, 1611.10300 & 2205.12827]

+more



A Family of Scrolls


...with cousins joining in...

Deform: $p_1(x, y) = x_0 y_0^5 + x_1 y_1^5 + x_2 y_0 y_1^4$ toric $F_{(4,1,0,\dots)}^{(n)}$

Now: $\mathfrak{s}_{1,1}(x, y) = \frac{x_0 y_0}{y_1^5} + \frac{x_2}{y_1^4} - \frac{x_1}{y_1^4}$ & $\mathfrak{s}_{1,2}(x, y) = \frac{x_0}{y_1} - \frac{x_2}{y_0} - \frac{x_1 y_1^4}{y_0^5}$

& $\det \left[\frac{\partial(p_1, \mathfrak{s}_{1,1}, \mathfrak{s}_{1,2}, x_3, \dots; y_0, y_1)}{\partial(x_0, x_1, x_2, x_3, \dots; y_0, y_1)} \right] = \text{const.}$

X_1	X_2	X_3	X_4	X_5	X_6	
1	1	1	1	0	0	$\leftarrow \mathbb{P}^4$
-4	-1	0	0	1	1	$\leftarrow \mathbb{P}^1$



Deform: $p_2(x, y) = x_0 y_0^5 + x_1 y_1^5 + x_2 y_0^2 y_1^3$ toric $F_{(3,2,0,\dots)}^{(n)}$

Now: $\mathfrak{s}_{2,1}(x, y) = \frac{x_0 y_0^2}{y_1^5} + \frac{x_2}{y_1^3} - \frac{x_1}{y_1^3}$ & $\mathfrak{s}_{2,2}(x, y) = \frac{x_0}{y_1^2} - \frac{x_2}{y_0^2} - \frac{x_1 y_1^3}{y_0^5}$


& $\det \left[\frac{\partial(p_2, \mathfrak{s}_{2,1}, \mathfrak{s}_{2,2}, x_3, \dots; y_0, y_1)}{\partial(x_0, x_1, x_2, x_3, \dots; y_0, y_1)} \right] = \text{const.}$

X_1	X_2	X_3	X_4	X_5	X_6	
1	1	1	1	0	0	$\leftarrow \mathbb{P}^4$
-3	-2	0	0	1	1	$\leftarrow \mathbb{P}^1$

... and $p_3(x, y) = x_0 y_0^5 + x_1 y_1^5 + x_2 y_0^2 y_1^3 + x_3 y_0^3 y_1^2$

$\rightarrow$ toric $F_{(2,2,1,\dots)}^{(n)}$

X_1	X_2	X_3	X_4	X_5	X_6	
1	1	1	1	0	0	$\leftarrow \mathbb{P}^4$
-2	-2	-1	0	1	1	$\leftarrow \mathbb{P}^1$



A Family of Scrolls



...with cousins joining in...

• Deform: $p_1(x, y) = x_0 y_0^5 + x_1 y_1^5 + x_2 y_0 y_1^4$ toric $F_{(4,1,0,\dots)}^{(n)}$

• Now: $\mathfrak{s}_{1,1}(x, y) = \frac{x_0 y_0}{y_1^5} + \frac{x_2}{y_1^4} - \frac{x_1}{y_1^4}$ & $\mathfrak{s}_{1,2}(x, y) = \frac{x_0}{y_1} - \frac{x_2}{y_0} - \frac{x_1 y_1^4}{y_0^5}$

• & $\det \left[\frac{\partial(p_1, \mathfrak{s}_{1,1}, \mathfrak{s}_{1,2}, x_3, \dots; y_0, y_1)}{\partial(x_0, x_1, x_2, x_3, \dots; y_0, y_1)} \right] = \text{const.}$

X_1	X_2	X_3	X_4	X_5	X_6	
1	1	1	1	0	0	$\leftarrow \mathbb{P}^4$
-4	-1	0	0	1	1	$\leftarrow \mathbb{P}^1$



[BH:1606.07420, 1611.10300 & 2205.12827]



+more

• Deform: $p_2(x, y) = x_0 y_0^5 + x_1 y_1^5 + x_2 y_0^2 y_1^3$ toric $F_{(3,2,0,\dots)}^{(n)}$

• Now: $\mathfrak{s}_{2,1}(x, y) = \frac{x_0 y_0^2}{y_1^5} + \frac{x_2}{y_1^3} - \frac{x_1}{y_1^3}$ & $\mathfrak{s}_{2,2}(x, y) = \frac{x_0}{y_1^2} - \frac{x_2}{y_0^2} - \frac{x_1 y_1^3}{y_0^5}$

• & $\det \left[\frac{\partial(p_2, \mathfrak{s}_{2,1}, \mathfrak{s}_{2,2}, x_3, \dots; y_0, y_1)}{\partial(x_0, x_1, x_2, x_3, \dots; y_0, y_1)} \right] = \text{const.}$

X_1	X_2	X_3	X_4	X_5	X_6	
1	1	1	1	0	0	$\leftarrow \mathbb{P}^4$
-3	-2	0	0	1	1	$\leftarrow \mathbb{P}^1$



• ... and $p_3(x, y) = x_0 y_0^5 + x_1 y_1^5 + x_2 y_0^2 y_1^3 + x_3 y_0^3 y_1^2$

• $\rightarrow$ toric $F_{(2,2,1,\dots)}^{(n)}$


for $n=3$,

$$F_{(2,2,1)}^{(3)} \approx F_{(1,1,0)}^{(3)}$$

[~Veronese]

Fano!


X_1	X_2	X_3	X_4	X_5	X_6	
1	1	1	1	0	0	$\leftarrow \mathbb{P}^4$
-2	-2	-1	0	1	1	$\leftarrow \mathbb{P}^1$



$F_{(1,1,0)}^{(3)}$

A Family of Scrolls


...in elaborate hierarchy



[BH:1606.07420, 1611.10300 & 2205.12827]

+more

Deg.	Biprojective Defining Section	Name, GLSM Charges															
	$p_0(x, y) = x_0 y_0^5 + x_1 y_1^5$	$F_5^{(3)}$															
$(-\frac{1}{5})$:	$\mathfrak{s}_0(x, y) = \left[\left(\frac{x_0}{y_1^5} - \frac{x_1}{y_0^5} \right) / p_0(x, y) \right]$	<table><tr><td>X_1</td><td>X_2</td><td>X_3</td><td>X_4</td><td>X_5</td></tr><tr><td>$-\frac{1}{5}$</td><td>1</td><td>1</td><td>0</td><td>0</td></tr><tr><td></td><td></td><td></td><td>1</td><td>1</td></tr></table>	X_1	X_2	X_3	X_4	X_5	$-\frac{1}{5}$	1	1	0	0				1	1
X_1	X_2	X_3	X_4	X_5													
$-\frac{1}{5}$	1	1	0	0													
			1	1													
	$p_1(x, y) = p_0(x, y) + x_2 y_0^4 y_1$	$F_{4,1}^{(3)}$															
$(-\frac{1}{2})$:	$\mathfrak{s}_{1a}(x, y) = \left[\left(\frac{x_0 y_0}{y_1^5} - \frac{x_1}{y_0^4} + \frac{x_2}{y_1^4} \right) / p_1(x, y) \right]$	<table><tr><td>X_1</td><td>X_2</td><td>X_3</td><td>X_4</td><td>X_5</td></tr><tr><td>$-\frac{1}{4}$</td><td>$-\frac{1}{1}$</td><td>1</td><td>0</td><td>0</td></tr><tr><td></td><td></td><td></td><td>1</td><td>1</td></tr></table>	X_1	X_2	X_3	X_4	X_5	$-\frac{1}{4}$	$-\frac{1}{1}$	1	0	0				1	1
X_1	X_2	X_3	X_4	X_5													
$-\frac{1}{4}$	$-\frac{1}{1}$	1	0	0													
			1	1													
$(-\frac{1}{2})$:	$\mathfrak{s}_{1b}(x, y) = \left[\left(\frac{x_0}{y_1} - \frac{x_1 y_1^4}{y_0^5} - \frac{x_2}{y_0} \right) / p_1(x, y) \right]$	<table><tr><td>X_1</td><td>X_2</td><td>X_3</td><td>X_4</td><td>X_5</td></tr><tr><td>$-\frac{1}{4}$</td><td>$-\frac{1}{1}$</td><td>1</td><td>0</td><td>0</td></tr><tr><td></td><td></td><td></td><td>1</td><td>1</td></tr></table>	X_1	X_2	X_3	X_4	X_5	$-\frac{1}{4}$	$-\frac{1}{1}$	1	0	0				1	1
X_1	X_2	X_3	X_4	X_5													
$-\frac{1}{4}$	$-\frac{1}{1}$	1	0	0													
			1	1													
	$p_2(x, y) = p_0(x, y) + x_2 y_0^3 y_1^2$	$F_{3,2}^{(3)}$															
$(-\frac{1}{3})$:	$\mathfrak{s}_{2a}(x, y) = \left[\left(\frac{x_0 y_0^2}{y_1^5} - \frac{x_1}{y_0^3} + \frac{x_2 y_0}{y_1^4} + \frac{x_3}{y_1^3} \right) / p_2(x, y) \right]$	<table><tr><td>X_1</td><td>X_2</td><td>X_3</td><td>X_4</td><td>X_5</td></tr><tr><td>$-\frac{1}{3}$</td><td>$-\frac{1}{2}$</td><td>1</td><td>0</td><td>0</td></tr><tr><td></td><td></td><td></td><td>1</td><td>1</td></tr></table>	X_1	X_2	X_3	X_4	X_5	$-\frac{1}{3}$	$-\frac{1}{2}$	1	0	0				1	1
X_1	X_2	X_3	X_4	X_5													
$-\frac{1}{3}$	$-\frac{1}{2}$	1	0	0													
			1	1													
$(-\frac{1}{1})$:	$\mathfrak{s}_{2b}(x, y) = \left[\left(\frac{x_0}{y_1^2} - \frac{x_1 y_1^3}{y_0^5} - \frac{x_2}{y_0^2} \right) / p_2(x, y) \right]$	<table><tr><td>X_1</td><td>X_2</td><td>X_3</td><td>X_4</td><td>X_5</td></tr><tr><td>$-\frac{1}{3}$</td><td>$-\frac{1}{2}$</td><td>1</td><td>0</td><td>0</td></tr><tr><td></td><td></td><td></td><td>1</td><td>1</td></tr></table>	X_1	X_2	X_3	X_4	X_5	$-\frac{1}{3}$	$-\frac{1}{2}$	1	0	0				1	1
X_1	X_2	X_3	X_4	X_5													
$-\frac{1}{3}$	$-\frac{1}{2}$	1	0	0													
			1	1													
	$p_3(x, y) = p_0(x, y) + x_2 y_0^4 y_1^1 + x_3 y_0^1 y_1^4$	$F_{3,1,1a}^{(3)} \approx_{\mathbb{R}} F_2^{(3)}$															
$(-\frac{1}{3})$:	$\mathfrak{s}_{3a}(x, y) = \left[\left(\frac{x_0 y_0}{y_1^4} - \frac{x_1 y_1}{y_0^4} + \frac{x_2}{y_1^3} - \frac{x_3}{y_0^3} \right) / p_3(x, y) \right]$	<table><tr><td>X_1</td><td>X_2</td><td>X_3</td><td>X_4</td><td>X_5</td></tr><tr><td>$-\frac{1}{3}$</td><td>$-\frac{1}{1}$</td><td>$-\frac{1}{1}$</td><td>0</td><td>0</td></tr><tr><td></td><td></td><td></td><td>1</td><td>1</td></tr></table>	X_1	X_2	X_3	X_4	X_5	$-\frac{1}{3}$	$-\frac{1}{1}$	$-\frac{1}{1}$	0	0				1	1
X_1	X_2	X_3	X_4	X_5													
$-\frac{1}{3}$	$-\frac{1}{1}$	$-\frac{1}{1}$	0	0													
			1	1													
$(-\frac{1}{1})$:	$\mathfrak{s}_{3b}(x, y) = \left[\left(\frac{x_0}{y_1} - \frac{x_1 y_1^4}{y_0^5} - \frac{x_2}{y_0} - \frac{x_3 y_1^3}{y_0^4} \right) / p_3(x, y) \right]$	<table><tr><td>X_1</td><td>X_2</td><td>X_3</td><td>X_4</td><td>X_5</td></tr><tr><td>$-\frac{1}{3}$</td><td>$-\frac{1}{1}$</td><td>$-\frac{1}{1}$</td><td>0</td><td>0</td></tr><tr><td></td><td></td><td></td><td>1</td><td>1</td></tr></table>	X_1	X_2	X_3	X_4	X_5	$-\frac{1}{3}$	$-\frac{1}{1}$	$-\frac{1}{1}$	0	0				1	1
X_1	X_2	X_3	X_4	X_5													
$-\frac{1}{3}$	$-\frac{1}{1}$	$-\frac{1}{1}$	0	0													
			1	1													
$(-\frac{1}{1})$:	$\mathfrak{s}_{3c}(x, y) = \left[\left(\frac{x_0 y_0^4}{y_1^5} - \frac{x_1}{y_0} + \frac{x_2 y_0^3}{y_1^4} + \frac{x_3}{y_1} \right) / p_3(x, y) \right]$	<table><tr><td>X_1</td><td>X_2</td><td>X_3</td><td>X_4</td><td>X_5</td></tr><tr><td>$-\frac{1}{3}$</td><td>$-\frac{1}{1}$</td><td>$-\frac{1}{1}$</td><td>0</td><td>0</td></tr><tr><td></td><td></td><td></td><td>1</td><td>1</td></tr></table>	X_1	X_2	X_3	X_4	X_5	$-\frac{1}{3}$	$-\frac{1}{1}$	$-\frac{1}{1}$	0	0				1	1
X_1	X_2	X_3	X_4	X_5													
$-\frac{1}{3}$	$-\frac{1}{1}$	$-\frac{1}{1}$	0	0													
			1	1													
	$p_4(x, y) = p_0(x, y) + x_2 y_0^4 y_1 + x_3 y_0^3 y_1^2$	$F_{3,1,1b}^{(3)} \approx_{\mathbb{R}} F_2^{(3)}$															
$(-\frac{1}{3})$:	$\mathfrak{s}_{4a}(x, y) = \left[\left(\frac{x_0 y_0^2}{y_1^5} - \frac{x_1}{y_0^3} + \frac{x_2 y_0}{y_1^4} + \frac{x_3}{y_1^3} \right) / p_4(x, y) \right]$	<table><tr><td>X_1</td><td>X_2</td><td>X_3</td><td>X_4</td><td>X_5</td></tr><tr><td>$-\frac{1}{3}$</td><td>$-\frac{1}{1}$</td><td>$-\frac{1}{1}$</td><td>0</td><td>0</td></tr><tr><td></td><td></td><td></td><td>1</td><td>1</td></tr></table>	X_1	X_2	X_3	X_4	X_5	$-\frac{1}{3}$	$-\frac{1}{1}$	$-\frac{1}{1}$	0	0				1	1
X_1	X_2	X_3	X_4	X_5													
$-\frac{1}{3}$	$-\frac{1}{1}$	$-\frac{1}{1}$	0	0													
			1	1													
$(-\frac{1}{1})$:	$\mathfrak{s}_{4b}(x, y) = \left[\left(\frac{x_0}{y_1} - \frac{x_1 y_1^4}{y_0^5} - \frac{x_2}{y_0} - \frac{x_3 y_1}{y_0^2} \right) / p_4(x, y) \right]$	<table><tr><td>X_1</td><td>X_2</td><td>X_3</td><td>X_4</td><td>X_5</td></tr><tr><td>$-\frac{1}{3}$</td><td>$-\frac{1}{1}$</td><td>$-\frac{1}{1}$</td><td>0</td><td>0</td></tr><tr><td></td><td></td><td></td><td>1</td><td>1</td></tr></table>	X_1	X_2	X_3	X_4	X_5	$-\frac{1}{3}$	$-\frac{1}{1}$	$-\frac{1}{1}$	0	0				1	1
X_1	X_2	X_3	X_4	X_5													
$-\frac{1}{3}$	$-\frac{1}{1}$	$-\frac{1}{1}$	0	0													
			1	1													
$(-\frac{1}{1})$:	$\mathfrak{s}_{4c}(x, y) = \left[\left(\frac{x_0 y_0}{y_1^2} - \frac{x_1 y_1^3}{y_0^4} + \frac{x_2}{y_1} - \frac{x_3}{y_0} \right) / p_4(x, y) \right]$	equivalent to $p_3(x, y)$															
	$p_5(x, y) = p_0(x, y) + x_2 y_0^3 y_1^2 + x_3 y_0^2 y_1^3$	$F_{2,2,1}^{(3)} \approx_{\mathbb{R}} F_{1,1}^{(3)}$															
$(-\frac{1}{2})$:	$\mathfrak{s}_{5a}(x, y) = \left[\left(\frac{x_0}{y_1^2} - \frac{x_1 y_1^3}{y_0^5} - \frac{x_2}{y_0^2} - \frac{x_3 y_1}{y_0^3} \right) / p_5(x, y) \right]$	<table><tr><td>X_1</td><td>X_2</td><td>X_3</td><td>X_4</td><td>X_5</td></tr><tr><td>$-\frac{1}{2}$</td><td>$-\frac{1}{2}$</td><td>$-\frac{1}{1}$</td><td>0</td><td>0</td></tr><tr><td></td><td></td><td></td><td>1</td><td>1</td></tr></table>	X_1	X_2	X_3	X_4	X_5	$-\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{1}$	0	0				1	1
X_1	X_2	X_3	X_4	X_5													
$-\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{1}$	0	0													
			1	1													
$(-\frac{1}{2})$:	$\mathfrak{s}_{5b}(x, y) = \left[\left(\frac{x_0 y_0^3}{y_1^5} - \frac{x_1}{y_0^2} + \frac{x_2 y_0}{y_1^3} + \frac{x_3}{y_1^2} \right) / p_5(x, y) \right]$	<table><tr><td>X_1</td><td>X_2</td><td>X_3</td><td>X_4</td><td>X_5</td></tr><tr><td>$-\frac{1}{2}$</td><td>$-\frac{1}{2}$</td><td>$-\frac{1}{1}$</td><td>0</td><td>0</td></tr><tr><td></td><td></td><td></td><td>1</td><td>1</td></tr></table>	X_1	X_2	X_3	X_4	X_5	$-\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{1}$	0	0				1	1
X_1	X_2	X_3	X_4	X_5													
$-\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{1}$	0	0													
			1	1													
$(-\frac{1}{1})$:	$\mathfrak{s}_{5c}(x, y) = \left[\left(\frac{x_0 y_0^2}{y_1^5} - \frac{x_1 y_1^2}{y_0^3} + \frac{x_2}{y_1} - \frac{x_3}{y_0} \right) / p_5(x, y) \right]$	<table><tr><td>X_1</td><td>X_2</td><td>X_3</td><td>X_4</td><td>X_5</td></tr><tr><td>$-\frac{1}{2}$</td><td>$-\frac{1}{2}$</td><td>$-\frac{1}{1}$</td><td>0</td><td>0</td></tr><tr><td></td><td></td><td></td><td>1</td><td>1</td></tr></table>	X_1	X_2	X_3	X_4	X_5	$-\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{1}$	0	0				1	1
X_1	X_2	X_3	X_4	X_5													
$-\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{1}$	0	0													
			1	1													





A Family of Scrolls

In a deformation reunion

- A deformation *family picture*:

[BH:1606.07420, 1611.10300 & 2205.12827]

+more





A Family of Scrolls

In a deformation reunion

- A deformation *family picture*:

[BH:1606.07420, 1611.10300 & 2205.12827]

+more





A Family of Scrolls

In a deformation reunion

- A deformation *family picture*:

[BH:1606.07420, 1611.10300 & 2205.12827]

+more



A Family of Scrolls

In a deformation reunion

- A deformation *family picture*:

- One complete intersection


- to “rule them all” (toric varieties)

- ...and CY hypersurfaces therein

- ...and a web of connected GLSMs

[BH:1606.07420, 1611.10300 & 2205.12827]

+more




A Family of Scrolls

...with combinatorial counterpoint

• A dihedral deformation *family picture*:

• 1951, Hirzebruch;
here, $m = 5$





A Family of Scrolls

...with combinatorial counterpoint

• A dihedral deformation *family picture*:

• 1951, Hirzebruch;
here, $m = 5$

• solid arrows:
explicit deformations,
by trial and directrix
identification



[BH:1606.07420,... 2605.06998]

A Family of Scrolls

...with combinatorial counterpoint

- A dihedral deformation *family picture*:

- 1951, Hirzebruch;
here, $m = 5$

- solid arrows:
explicit deformations,
by trial and directrix
identification

- dashed arrows:
non-algebraic map
(Wall “mod m ” diffeo;
(...perhaps...)
~Veronese-twisted,
smooth-deformed)


$$F_{2,1,1,0}^{(4)} \rightsquigarrow F_{0,0,0,0}^{(4)}$$

~ Gross [alg-geom/9402014]

& Ruan [DMJ 83.2 (1996) 461]

~ Wall & Veronese: $\vec{m} \rightsquigarrow \vec{m} \bmod n$

$$\left[\begin{array}{c|c} \mathbb{P}^n & 1 \\ \hline \mathbb{P}^1 & m \end{array} \right]$$




$$F_{3,3}^{(3)} \quad F_{4,3}^{(3)} \quad F_{5,3}^{(3)} \quad F_{6,3}^{(3)} \quad F_{7,3}^{(3)} \quad F_{8,3}^{(3)} \quad \dots$$

$$F_{2,2}^{(3)} \quad F_{3,2}^{(3)} \quad F_{4,2}^{(3)} \quad F_{5,2}^{(3)} \quad F_{6,2}^{(3)} \quad F_{7,2}^{(3)} \quad \dots$$

$$F_{1,1}^{(3)} \quad F_{2,1}^{(3)} \quad F_{3,1}^{(3)} \quad F_{4,1}^{(3)} \quad F_{5,1}^{(3)} \quad F_{6,1}^{(3)} \quad \dots$$

$$F_{0,0}^{(3)} \quad F_{1,0}^{(3)} \quad F_{2,0}^{(3)} \quad F_{3,0}^{(3)} \quad F_{4,0}^{(3)} \quad F_{5,0}^{(3)} \quad \dots \quad m_1$$

$$F_{3,1,1}^{(n)} \rightsquigarrow F_{2,0,0}^{(n)}$$





[BH:1606.07420,... 2605.06998]

and Their Offspring

...back to the main motif 

- CY: $F_m^{(n)}[c_1]$, i.e., $F_m^{(n)} \in \left[\mathbb{P}^n \middle| \begin{matrix} 1 \\ m \end{matrix} \middle| \begin{matrix} n \\ 2-m \end{matrix} \right]$, i.e., need $\deg = \binom{n}{2-m}$
- $f_0(X) = \Pi X := (X_1 = \mathfrak{z}(x, y)) \cdots (X_n = x_n)(X_{n+1} = y_0)(X_{n+2} = y_1) = \text{“fundamental monom.”}$
 - is not transverse & $\{\Pi X = 0\}$ is singular (“ $(n+1)$ -horon”)






and Their Offspring

...back to the main motif 


- CY: $F_m^{(n)}[c_1]$, i.e., $F_m^{(n)} \in \left[\begin{array}{c|c} \mathbb{P}^n & 1 \\ \hline \mathbb{P}^1 & m \end{array} \middle| \begin{array}{c} n \\ 2-m \end{array} \right]$, i.e., need $\deg = \binom{n}{2-m}$
- $f_0(X) = \Pi X := (X_1 = \mathfrak{s}(x, y)) \cdots (X_n = x_n)(X_{n+1} = y_0)(X_{n+2} = y_1) =$ “fundamental monom.”
 - is not transverse & $\{\Pi X = 0\}$ is singular (“ $(n+1)$ -horon”)
- Deform: $f_0(X) = \Pi X \rightsquigarrow \Pi X + \epsilon^i \delta_i \Pi X$, $\delta_i \Pi X := \mathfrak{m}_i(X)(\partial_i \Pi X)$
 where $\deg[\mathfrak{m}_i(X)] = \deg[X_i]$ & $\partial_i \mathfrak{m}_i(X) = 0$ ($\sim$ Demazure roots)



and Their Offspring


...back to the main motif 


- CY: $F_m^{(n)}[c_1]$, i.e., $F_m^{(n)} \in \left[\mathbb{P}^n \middle| \begin{smallmatrix} 1 \\ m \end{smallmatrix} \middle| \begin{smallmatrix} n \\ 2-m \end{smallmatrix} \right]$, i.e., need $\deg = \binom{n}{2-m}$
- $f_0(X) = \Pi X := (X_1 = \mathfrak{z}(x, y)) \cdots (X_n = x_n)(X_{n+1} = y_0)(X_{n+2} = y_1) = \text{“fundamental monom.”}$
 - is not transverse & $\{\Pi X = 0\}$ is singular (“ $(n+1)$ -horon”)
- Deform: $f_0(X) = \Pi X \rightsquigarrow \Pi X + \epsilon^i \delta_i \Pi X$, $\delta_i \Pi X := \mathfrak{m}_i(X)(\partial_i \Pi X)$
 where $\deg[\mathfrak{m}_i(X)] = \deg[X_i]$ & $\partial_i \mathfrak{m}_i(X) = 0$ ($\sim$ Demazure roots)
 - in QM, $|n\rangle^{(1)} = \sum_{m \neq n} c_m |m\rangle^{(0)}$; also, $X_i = 0 \not\Rightarrow \delta_i \Pi X = 0$
 - $(\delta_i \Pi X) \cap (\delta_j \Pi X)$ are X_i - and X_j -independent and delimit $[\delta_i \Pi X]$ & $[\delta_j \Pi X]$
 so $\langle X_i \rangle, \langle X_j \rangle = 0 \not\Rightarrow \langle [\delta_i \Pi X] \cap [\delta_j \Pi X] \rangle = 0$; etc. **they form a poset**



and Their Offspring

...back to the main motif 






and Their Offspring

...back to the main motif 

- CY: $F_m^{(n)}[c_1]$, i.e., $F_m^{(n)} \in \left[\mathbb{P}^n \middle| \begin{matrix} 1 \\ m \end{matrix} \middle| \begin{matrix} n \\ 2-m \end{matrix} \right]$, i.e., need $\deg = \binom{n}{2-m}$
- $f_0(X) = \Pi X := (X_1 = \mathfrak{z}(x, y)) \cdots (X_n = x_n)(X_{n+1} = y_0)(X_{n+2} = y_1) = \text{“fundamental monom.”}$
 - is not transverse & $\{\Pi X = 0\}$ is singular (“ $(n+1)$ -horon”)
- Deform: $f_0(X) = \Pi X \rightsquigarrow \Pi X + \epsilon^i \delta_i \Pi X$, $\delta_i \Pi X := \mathfrak{m}_i(X)(\partial_i \Pi X)$
 where $\deg[\mathfrak{m}_i(X)] = \deg[X_i]$ & $\partial_i \mathfrak{m}_i(X) = 0$ ($\sim$ Demazure roots)
 - in QM, $|n\rangle^{(1)} = \sum_{m \neq n} c_m |m\rangle^{(0)}$; also, $X_i = 0 \not\Rightarrow \delta_i \Pi X = 0$
 - $(\delta_i \Pi X) \cap (\delta_j \Pi X)$ are X_i - and X_j -independent and delimit $[\delta_i \Pi X]$ & $[\delta_j \Pi X]$
 so $\langle X_i \rangle, \langle X_j \rangle = 0 \not\Rightarrow \langle [\delta_i \Pi X] \cap [\delta_j \Pi X] \rangle = 0$; etc. **they form a poset**
- Combinatorially, $\Pi X \sqcup \bigsqcup_i [\delta_i \Pi X] \stackrel{\nabla}{\simeq} \{\Pi X = 0 \mid \text{S.R., ...}\}$
- Also, $\partial_i \perp \delta_i \Pi X$ & generate a dual poset




and Their Offspring

(bird's eye view)



transpolar (face-wise polar)
images of fan-spanning polytopes




Monomial collections with $U(1)^2$ -charges of ΠX form posets, with “deformation fronts” = facets

The shape changes throughout the ϵ_{ik} -space indicate that — *convexity is overrated*

In all deformed models, there exist transverse *regular* polynomials (*nonnegative exponents*)...

except in $F_5^{(3)}[c_1]$: requires **rational monomials** to be transverse, & (L'Hôpital/intrinsic limit) smooth



...in the Mirror

& Non-Convex Mirrors

$m=3$ —2D Proof-of-Concept—

$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m}$$

•

•

•

•

•


•

•

•

•

universal
 $X_1 X_2 X_3 X_4$





...in the Mirror

& Non-Convex Mirrors

$m=3$ —2D Proof-of-Concept—

$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m} \oplus X_1^0 X_2^2 (X_3 \oplus X_4)^{2-1m}$$







...in the Mirror

& Non-Convex Mirrors

$m=3$ —2D Proof-of-Concept—

$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m} \oplus X_1^0 X_2^2 (X_3 \oplus X_4)^{2-1m}$$





...in the Mirror

& Non-Convex Mirrors


$m=3$ —2D Proof-of-Concept—


$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m} \oplus X_1^0 X_2^2 (X_3 \oplus X_4)^{2-1m}$$

Transpolar ($\approx$ dual):

$\Delta \rightarrow \bigsqcup_i (\text{convex } \Theta_i)$;

Compute $\Theta_i \rightarrow \Theta_i^\circ := \{v: \langle v | \forall u \in \Theta_i \rangle + 1 > 0\}$





...in the Mirror

& Non-Convex Mirrors


$m=3$ —2D Proof-of-Concept—

$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m} \oplus X_1^0 X_2^2 (X_3 \oplus X_4)^{2-1m}$$

• Transpolar ($\approx$ dual):

• $\Delta \rightarrow \bigsqcup_i (\text{convex } \Theta_i)$;

• Compute $\Theta_i \rightarrow \Theta_i^\circ := \{v: \langle v | \forall u \in \Theta_i \rangle + 1 > 0\}$



•

•

•


•


•

•

•

•





...in the Mirror

& Non-Convex Mirrors


$m=3$ —2D Proof-of-Concept—

$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m} \oplus X_1^0 X_2^2 (X_3 \oplus X_4)^{2-1m}$$

• Transpolar ($\approx$ dual):

• $\Delta \rightarrow \bigsqcup_i (\text{convex } \Theta_i)$;

• Compute $\Theta_i \rightarrow \Theta_i^\circ := \{v: \langle v | \forall u \in \Theta_i \rangle + 1 > 0\}$



•


•


•

•

•

•





...in the Mirror

& Non-Convex Mirrors



$m=3$ —2D Proof-of-Concept—


$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m} \oplus X_1^0 X_2^2 (X_3 \oplus X_4)^{2-1m}$$

• Transpolar ($\approx$ dual):

• $\Delta \rightarrow \bigsqcup_i (\text{convex } \Theta_i)$;

• Compute $\Theta_i \rightarrow \Theta_i^\circ := \{v: \langle v | \forall u \in \Theta_i \rangle + 1 > 0\}$





...in the Mirror

& Non-Convex Mirrors


$m=3$ —2D Proof-of-Concept—

$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m} \oplus X_1^0 X_2^2 (X_3 \oplus X_4)^{2-1m}$$

Transpolar ($\approx$ dual):


$\Delta \rightarrow \bigsqcup_i (\text{convex } \Theta_i)$;

Compute $\Theta_i \rightarrow \Theta_i^\circ := \{v: \langle v | \forall u \in \Theta_i \rangle + 1 > 0\}$



universal
 $X_1 X_2 X_3 X_4$





...in the Mirror

& Non-Convex Mirrors


$m=3$ —2D Proof-of-Concept—

$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m} \oplus X_1^0 X_2^2 (X_3 \oplus X_4)^{2-1m}$$

• Transpolar ($\approx$ dual):


• $\Delta \rightarrow \bigsqcup_i (\text{convex } \Theta_i)$;

• Compute $\Theta_i \rightarrow \Theta_i^\circ := \{v: \langle v | \forall u \in \Theta_i \rangle + 1 > 0\}$



universal
 $X_1 X_2 X_3 X_4$





...in the Mirror

& Non-Convex Mirrors


$m=3$ —2D Proof-of-Concept—

$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m} \oplus X_1^0 X_2^2 (X_3 \oplus X_4)^{2-1m}$$

• Transpolar ($\approx$ dual):

• $\Delta \rightarrow \bigsqcup_i (\text{convex } \Theta_i)$;

• Compute $\Theta_i \rightarrow \Theta_i^\circ := \{v: \langle v | \forall u \in \Theta_i \rangle + 1 > 0\}$



•

•

•

•

•

•

•

•

•


•

•

universal
 $X_1 X_2 X_3 X_4$



•



...in the Mirror

& Non-Convex Mirrors


$m=3$ —2D Proof-of-Concept—

$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m} \oplus X_1^0 X_2^2 (X_3 \oplus X_4)^{2-1m}$$

• Transpolar ($\approx$ dual):

• $\Delta \rightarrow \bigsqcup_i (\text{convex } \Theta_i)$;

• Compute $\Theta_i \rightarrow \Theta_i^\circ := \{v: \langle v | \forall u \in \Theta_i \rangle + 1 > 0\}$



•

•

•

•

•

•

•


•

•

universal
 $X_1 X_2 X_3 X_4$

•

•



...in the Mirror

& Non-Convex Mirrors

$m=3$ —2D Proof-of-Concept—


$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m} \oplus X_1^0 X_2^2 (X_3 \oplus X_4)^{2-1m}$$


Transpolar ($\approx$ dual):

$\Delta \rightarrow \bigsqcup_i (\text{convex } \Theta_i)$;

Compute $\Theta_i \rightarrow \Theta_i^\circ := \{v: \langle v | \forall u \in \Theta_i \rangle + 1 > 0\}$

(Re)assemble dually
 $(\theta_i \cap \theta_j)^\circ = [\theta_i^\circ, \theta_j^\circ]$
 with “neighbors”





...in the Mirror

& Non-Convex Mirrors

$m=3$ —2D Proof-of-Concept—

$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m} \oplus X_1^0 X_2^2 (X_3 \oplus X_4)^{2-1m}$$


Transpolar ($\approx$ dual):

$\Delta \rightarrow \bigsqcup_i (\text{convex } \Theta_i)$;

Compute $\Theta_i \rightarrow \Theta_i^\circ := \{v: \langle v | \forall u \in \Theta_i \rangle + 1 > 0\}$

(Re)assemble dually
 $(\theta_i \cap \theta_j)^\circ = [\theta_i^\circ, \theta_j^\circ]$
 with “neighbors”

Swap roles
 $X_i \leftrightarrow Y_i$



$$\mathfrak{G}_{ij} = \langle u_i, v_j \rangle + 1$$

$u_i \in \Delta_X^\star, v_j \in \Delta_X$
 mirror = transposition

$$F_m[c_1] \xleftrightarrow{\text{MM}} \nabla F_m[c_1]$$



...in the Mirror

& Non-Convex Mirrors

$m=3$ —2D Proof-of-Concept—

$$X_1^2 X_2^0 (X_3 \oplus X_4)^{2+1m} \oplus X_1^1 X_2^1 (X_3 \oplus X_4)^{2+0m} \oplus X_1^0 X_2^2 (X_3 \oplus X_4)^{2-1m}$$

Transpolar ($\approx$ dual):

$\Delta \rightarrow \bigsqcup_i (\text{convex } \Theta_i)$;

Compute $\Theta_i \rightarrow \Theta_i^\circ := \{v: \langle v | \forall u \in \Theta_i \rangle + 1 > 0\}$

(Re)assemble dually
 $(\theta_i \cap \theta_j)^\circ = [\theta_i^\circ, \theta_j^\circ]$
 with “neighbors”

Swap roles
 $X_i \leftrightarrow Y_i$

$$f(X) = X_1^2 X_3^5 + X_1^2 X_4^5 + X_2^2 X_4^{-1} + X_2^2 X_3^{-1}$$

$$f^\Gamma(Y) = Y_1^2 Y_2^2 + Y_3^2 Y_4^2 + Y_1^5 Y_4^{-1} + Y_2^5 Y_3^{-1}$$

$$\mathfrak{E} = \begin{bmatrix} 2 & 0 & 5 & 0 \\ 2 & 0 & 0 & 5 \\ 0 & 2 & 0 & -1 \\ 0 & 2 & -1 & 0 \end{bmatrix}$$

$$\mathfrak{E}_{ij} = \langle u_i, v_j \rangle + 1$$

$$u_i \in \Delta_X^\star, v_j \in \Delta_X$$

mirror = transposition

$\text{rk}[\mathfrak{E}] = 3$, so need to
 chisel both $\Delta_{F_3}^\star$ and

also $\Delta_{F_3} =: \Delta_{\nabla F_3}^\star$
 to 0-enclosing
 simplices

Who's
 $\nabla F_m^{(n)}?! \dots$

(pre)complex
 algebraic
 geometry


conVEX
 polytopes


$$F_m[c_1] \xleftrightarrow{\text{MM}} \nabla F_m[c_1]$$

(pre)symplectic geometry

universal
 $X_1 X_2 X_3 X_4$

'92: Khovanskii
 + Pukhlikov
 '93: Karshon
 + Tolman
 '99: Hattori
 + Masuda
 + lots of





...in the Mirror

& Non-Convex Mirrors

$m=3$ —2D Proof-of-Concept—

What of self-crossing multifans & multitoes that span them?

Good question!

“sandwiched” between toric varieties (far right)

and (unitary) torus manifolds (far left) with $(S^1)^n$ -equivariant almost complex structure


$\mathbb{C}^* \approx S^1 \times \mathbb{R}_+ =$ “circular” & “radial” action

Topological Torus Manifolds? [Ishida, Fukukawa & Masuda, 2013]

Top cone orientation $\leftrightarrow$ rel. orientation btw. M & $\text{cpx}[T_M \otimes \mathbb{R}^{2k}]$...

They have cohomology rings generated by Cox-like coords & “ $\mathcal{K}_M^*$ ”

Guess: *obverse top-cones* obstruct global almost complex structure (...& break supersymmetry...?)




top cones encode characteristic points; obverse top cones encode points where complex structure “breaks”

...in the Mirror

Summary 

—Proof-of-Concept—

- CY($n-1$)-folds in Hirzebruch n -folds, $X_m^{(n-1)} \in F_m^{(n)}[c_1]$



...in the Mirror

Summary

—Proof-of-Concept—


- CY($n-1$)-folds in Hirzebruch n -folds, $X_m^{(n-1)} \in F_m^{(n)}[c_1]$

Each distinct $F_{\vec{m}}^{(n)}$ harbors a combinatorial web of transposition mirror models

In generic $F_{\vec{m}}^{(n)}$, the ΠX has plenty of regular smoothing deformations (with only nonnegative exponents)

No regular smoothing deformations remain in the central model: only rational monomials remain

All these distinct toric varieties are found within the same explicit deformation family




How Hard Can it Be?

Constructing $CY \subset$ Some “Nice” Ambient Space

- Reduce to 0 dimensions: $\mathbb{P}^4[5] \rightarrow \mathbb{P}^3[4] \rightarrow \mathbb{P}^2[3] \rightarrow \mathbb{P}^1[2]$

1611.10300
2205.12827
2403.07139
2501.11684
2502.08002
2605.06998
2605.24724
to be cont'd








Just add
some dimensions



Counting with multiplicity
 $\mathcal{X} \in A[c_1]$

Eldorado Ho!



Thank You!

1611.10300
2205.12827
2403.07139
2501.11684
2502.08002
2605.06998
2605.24724
to be cont'd

Eldorado Ho!